

A NEW EXACT PENALTY FUNCTION

WALTRAUD HUYER[†] AND ARNOLD NEUMAIER[†]

Abstract. For constrained smooth or nonsmooth optimization problems, new continuously differentiable penalty functions are derived. They are proved exact in the sense that under some nondegeneracy assumption, local optimizers of a nonlinear program are precisely the optimizers of the associated penalty function. This is achieved by augmenting the dimension of the program by a variable that controls both the weight of the penalty terms and the regularization of the nonsmooth terms.

Key words. constrained optimization, nonlinear programming, nonsmooth optimization, exact penalty function, Mangasarian–Fromovitz condition, augmented Lagrangian

AMS subject classification. 90C30

1. Introduction. Smooth nonlinear programs are traditionally solved by augmenting the objective function or a corresponding Lagrangian function using penalty or barrier terms to take account of the constraints (see, e.g., the surveys [3, 15]). The resulting merit function is then optimized using either standard unconstrained (or bound constrained) optimization software or sequential quadratic programming (SQP) techniques. Independently of the technique used, the merit function always depends on a small parameter ε (or a large parameter $\rho = \varepsilon^{-1}$); as $\varepsilon \rightarrow 0$, minimizers of the merit function converge to the set of minimizers of the original problem. In some SQP approaches, one uses instead so-called exact penalty functions that produce exact optimizers already at sufficiently small positive values of ε . In return, these exact penalty functions have the disadvantage that the evaluation of the merit function either needs Jacobian information (to estimate multipliers), or (for l_1 or l_∞ penalties) it is no longer smooth. In addition, both kinds of penalty functions may be unbounded below even when the constrained problem is bounded, which may make it difficult or impossible to locate a minimizer.

For nonsmooth nonlinear programs, solution techniques are much less developed, and often restricted to the convex or unconstrained case; in the latter case, constraints are usually handled by an l_∞ exact penalty function (e.g. in SOLVOPT [6]). The various approaches are based on combinations of subgradient methods (e.g., [13, 4]), Moreau–Yosida regularization (e.g., [8, 9, 14]), or bundle techniques (e.g., [5, 7, 10]). The regularization again depends on a small smoothing parameter $\varepsilon > 0$ such that for $\varepsilon \rightarrow 0$ the original nonsmooth functions are recovered.

In the following, we discuss a new merit function for smooth or nonsmooth optimization problems with equality, inequality and bound constraints that

- has good smoothness and exactness properties,
- remains bounded below under reasonable conditions,
- combines regularization with penalty techniques, and
- is flexible enough to give enough freedom for incorporating available Lagrange multiplier estimates.

The most important new idea is that the merit function is considered as a function of x and ε simultaneously, with the property that under appropriate assumptions, the

[†]Institut für Mathematik, Universität Wien, Strudlhofgasse 4, A-1090 Wien, Austria
(Waltraud.Huyer@univie.ac.at, Arnold.Neumaier@univie.ac.at).

minimizer (x^*, ε^*) of the merit function satisfies $\varepsilon^* = 0$, so that x^* solves the original problem.

This paper is organized as follows. In §2, the case of a smooth constrained optimization problem with equality constraints and bound constraints is considered. A penalty function is introduced and it is proved that under certain assumptions, all local minimizers of this penalty function have the form (x, ε) with $\varepsilon = 0$ and x a solution of the original problem. A converse of this can be proved in much greater generality, namely for nonsmooth functions that are suitably regularized. Therefore, in §3, a nonsmooth objective function and nonsmooth constraints are replaced by regularized functions and regularization recipes for some common nonsmooth functions are given. As a preparation for the exactness proof, §4 proves some results involving regular zeros of a not necessarily smooth function. In §5 the penalty function is generalized to the regularized problem and it is proved that, for a solution x^* of the constrained optimization problem, $(x^*, 0)$ is a minimizer of the penalty function. In §6 we illustrate our theory with an example, where the traditional penalty functions are unbounded. Finally, in §7 our penalty function is generalized to problems involving in addition inequality constraints; analogous results to those for equality constraints are shown to hold by reducing this case to the previous one with the aid of slack variables.

Notation. In the following, the absolute value of a vector is defined componentwise, $|x| := (|x_1|, \dots, |x_n|)^T$. Similarly, vector inequalities are understood componentwise. The norm used throughout is the Euclidean norm $\|x\| = \sqrt{\sum x_k^2}$, and $B[x_0; r]$ denotes a closed Euclidean ball around x_0 with radius r . The subvector of x indexed by the indices in J is denoted by x_J , and $A_{\cdot J}$ denotes the matrix consisting of the columns of a matrix A indexed by the indices in J . Sets of the form

$$\mathbf{x} = [\underline{x}, \bar{x}] := \{x \in \mathbb{R}^n \mid \underline{x} \leq x \leq \bar{x}\},$$

where the *lower bound* $\underline{x} \in (\mathbb{R} \cup \{-\infty\})^n$ and the *upper bound* $\bar{x} \in (\mathbb{R} \cup \{\infty\})^n$ are vectors containing proper or infinite bounds on the components of x and $\underline{x} \leq \bar{x}$, are referred to as *n-dimensional boxes*.

2. The smooth case. In this section we propose a class of penalty functions for the smooth constrained nonlinear optimization problem

$$(2.1) \quad \begin{array}{ll} \min & f(x) \\ \text{s.t.} & x \in [u, v], \quad F(x) = 0, \end{array}$$

where $[u, v]$ is a box in \mathbb{R}^n with nonempty interior, $f : D \rightarrow \mathbb{R}$ and $F : D \rightarrow \mathbb{R}^m$ are continuously differentiable in an open set D containing $[u, v]$. We fix $w \in \mathbb{R}^m$ and consider the equivalent problem

$$(2.2) \quad \begin{array}{ll} \min & f(x) \\ \text{s.t.} & F_i(x) = \varepsilon w_i \quad (i = 1, \dots, m), \\ & x \in [u, v], \quad \varepsilon = 0. \end{array}$$

This motivates the definition of the penalty function f_σ on $D \times [0, \bar{\varepsilon}]$ by

$$(2.3) \quad f_\sigma(x, \varepsilon) = \begin{cases} f(x) & \text{if } \varepsilon = \Delta(x, \varepsilon) = 0, \\ f(x) + \frac{1}{2\varepsilon} \cdot \frac{\Delta(x, \varepsilon)}{1 - q\Delta(x, \varepsilon)} + \sigma\beta(\varepsilon) & \text{if } \varepsilon > 0, \Delta(x, \varepsilon) < q^{-1}, \\ \infty & \text{otherwise,} \end{cases}$$

with the *constraint violation measure*

$$(2.4) \quad \Delta(x, \varepsilon) := \|\varepsilon w - F(x)\|^2,$$

where, in addition, $\bar{\varepsilon} > 0$ and $q > 0$ are fixed and $\beta : [0, \bar{\varepsilon}] \rightarrow [0, \infty)$ is continuous and continuously differentiable on $(0, \bar{\varepsilon}]$ with $\beta(0) = 0$. The surrogate optimization problem then reads

$$(2.5) \quad \begin{array}{ll} \min & f_\sigma(x, \varepsilon) \\ \text{s.t.} & (x, \varepsilon) \in [u, v] \times [0, \bar{\varepsilon}]. \end{array}$$

Note that $f_\sigma(x, \varepsilon)$ is continuously differentiable in

$$D_\sigma = \{(x, \varepsilon) \in D \times (0, \bar{\varepsilon}) \mid \Delta(x, \varepsilon) < q^{-1}\},$$

with continuous limits at the part of the boundary where the limit values are finite, in particular at $(x, 0)$ with feasible x . Moreover,

$$(2.6) \quad f_\sigma(x, \varepsilon) = f(x) + \frac{\varepsilon}{2} \cdot \frac{\|w\|^2}{1 - q\|w\|^2\varepsilon^2} + \sigma\beta(\varepsilon) \geq f(x) = f_\sigma(x, 0) \quad \text{if } F(x) = 0.$$

The shift by εw in the definition of the constraint violation measure allows one to incorporate Lagrange multiplier estimates (that serve to be able to work with better conditioned Hessians; see the remark at the end of this section).

The denominator $1 - q\Delta(x, \varepsilon)$ is included since it forces the level sets of f_σ to remain in the set $\{(x, \varepsilon) \in \mathbb{R}^n \mid \Delta(x, \varepsilon) < q^{-1}\}$ and hence in some sense close to the feasible set of (2.1). In particular, in many cases where the traditional quadratic penalty function (where $w = 0$ and $q = 0$) is unbounded below, moderate positive values for q give a well-behaved penalty problem (cf. the example in §6). Indeed, f_σ is bounded below on $[u, v] \times [0, \bar{\varepsilon}]$ whenever $f(x)$ is bounded below on the set

$$(2.7) \quad D' = \{x \in [u, v] \mid \|F(x)\| \leq q^{-1/2} + \bar{\varepsilon}\|w\|\}.$$

This is a reasonable condition since it always holds when f is bounded below on the feasible set, $\bar{\varepsilon}$ is small enough, and q is large enough.

The term $\sigma\beta(\varepsilon)$ is included since it allows to optimize simultaneously on x and ε , thus automatizing the adaptation of the penalty factor $1/2\varepsilon$. Intuitively, many slices with different, fixed values of ε that are optimized in traditional quadratic penalty methods are arranged consecutively, and translated by the term $\sigma\beta(\varepsilon)$ in such a way that the minimizers form a curve with decreasing function values as $\varepsilon \rightarrow 0$. Therefore, simultaneous optimization over both x and ε automatically leads to a local minimum at $\varepsilon = 0$. Of course, we need conditions guaranteeing that the term $\sigma\beta(\varepsilon)$ is enough to cause this behavior of the penalty function. As we shall see in §5, $\beta(\varepsilon) = \sqrt{\varepsilon}$ is an appropriate (but not the only possible) choice.

We say that the *Mangasarian–Fromovitz condition* (see [11]) for (2.1) holds at $x \in [u, v]$ if $F'(x)$ has full rank and there is a $p \in \mathbb{R}^n$ with $F'(x)p = 0$ and

$$p_i \begin{cases} > 0 & \text{if } x_i = u_i, \\ < 0 & \text{if } x_i = v_i. \end{cases}$$

THEOREM 2.1. *In addition to the general assumptions mentioned after (2.1) and after (2.4), assume that the set (2.7) is bounded, that each $x \in D'$ satisfies the Mangasarian–Fromovitz condition, and that*

$$(2.8) \quad \beta'(\varepsilon) \geq \beta_1 > 0 \quad \text{for } 0 < \varepsilon < \bar{\varepsilon}.$$

If σ is sufficiently large, there is no Kuhn–Tucker point (x, ε) of (2.5) with $\varepsilon > 0$.

In particular, for sufficiently large σ , every local minimizer (x^*, ε^*) of the penalty problem (2.5) with finite $f_\sigma(x^*, \varepsilon^*)$ has the form $(x^*, 0)$, where x^* is a local minimizer of the original problem (2.1).

Proof. If (x, ε) is a Kuhn–Tucker point of (2.5) with $\varepsilon > 0$, then there exist vectors $y, z \in \mathbb{R}^{n+1}$ such that

$$\begin{aligned} \nabla f_\sigma(x, \varepsilon) &= y - z, \\ \inf(y_i, x_i - u_i) &= \inf(z_i, v_i - x_i) = 0, \quad i = 1, \dots, n, \\ y_{n+1} &= \inf(z_{n+1}, \bar{\varepsilon} - \varepsilon) = 0, \end{aligned}$$

where $\nabla f_\sigma(x, \varepsilon)$ is the gradient of f_σ with respect to (x, ε) . The assertion of the theorem is proved by contradiction. Assume that there exists a sequence $(x^k, \varepsilon_k, \sigma_k)$, $\varepsilon_k \neq 0$ for all k , $\sigma_k \rightarrow \infty$ as $k \rightarrow \infty$, where (x^k, ε_k) is a Kuhn–Tucker point of f_{σ_k} . We use the abbreviation $\Delta_k := \Delta(x^k, \varepsilon_k)$. The point x^k satisfies

$$\|F(x^k)\| \leq \Delta_k^{1/2} + \varepsilon_k \|w\| \leq q^{-1/2} + \bar{\varepsilon} \|w\|;$$

hence $x^k \in D'$. Since D' is closed and bounded, we may restrict to a subsequence if necessary, and assume that

$$(2.9) \quad \lim_{k \rightarrow \infty} \varepsilon_k = \varepsilon^* \in [0, \bar{\varepsilon}] \quad \text{and} \quad \lim_{k \rightarrow \infty} x^k = x^* \in D'.$$

The condition $\frac{\partial}{\partial \varepsilon} f_{\sigma_k}(x^k, \varepsilon_k) \leq 0$ yields

$$(2.10) \quad q\Delta_k^2 + \varepsilon_k^2 \|w\|^2 + 2\varepsilon_k^2 (1 - q\Delta_k)^2 \sigma_k \beta'(\varepsilon_k) \leq \|F(x^k)\|^2,$$

with equality in the case $\varepsilon_k \neq \bar{\varepsilon}$. Since the right-hand side is bounded and $\sigma_k \rightarrow \infty$, this yields (in view of (2.8) and (2.9))

$$(2.11) \quad \varepsilon^* = 0 \quad \text{or} \quad \Delta^* = q^{-1},$$

where $\Delta^* := \Delta(x^*, \varepsilon^*)$. The derivatives with respect to x give

$$(2.12) \quad f_{x_i}(x^k) + \frac{1}{(1 - q\Delta_k)^2 \varepsilon_k} (F'(x^k)^T (F(x^k) - \varepsilon_k w))_i \begin{cases} \geq 0 & \text{if } x_i^k = u_i, \\ = 0 & \text{if } u_i < x_i^k < v_i, \\ \leq 0 & \text{if } x_i^k = v_i, \end{cases}$$

or

$$(F'(x^k)^T (F(x^k) - \varepsilon_k w))_i + (1 - q\Delta_k)^2 \varepsilon_k f_{x_i}(x^k) \begin{cases} \geq 0 & \text{if } x_i^k = u_i, \\ = 0 & \text{if } u_i < x_i^k < v_i, \\ \leq 0 & \text{if } x_i^k = v_i, \end{cases}$$

where f_{x_i} denotes the partial derivative of f with respect to x_i . By passing to the limit, using (2.9) and (2.11), we obtain

$$(2.13) \quad (F'(x^*)^T (F(x^*) - \varepsilon^* w))_i \begin{cases} \geq 0 & \text{if } x_i^* = u_i, \\ = 0 & \text{if } u_i < x_i^* < v_i, \\ \leq 0 & \text{if } x_i^* = v_i. \end{cases}$$

Since $x^* \in D'$, the Mangasarian–Fromovitz condition holds for $x = x^*$ and some vector $p \in \mathbb{R}^n$. Let $I_1 := \{i \mid x_i^* = u_i\}$, $I_2 := \{i \mid x_i^* = v_i\}$ and $w^* := F(x^*) - \varepsilon^* w$. Then

$$0 = (F'(x^*)p)^T w^* = \sum_{i \in I_1} p_i (F'(x^*)^T w^*)_i + \sum_{i \in I_2} p_i (F'(x^*)^T w^*)_i,$$

and the Mangasarian–Fromovitz condition and (2.13) imply $(F'(x^*)^T w^*)_i = 0$ for $i \in I_1 \cup I_2$ and thus $F'(x^*)^T w^* = 0$. Now the fact that $F'(x^*)$ has full rank yields $w^* = 0$, giving

$$(2.14) \quad F(x^*) - \varepsilon^* w = 0.$$

Hence $\Delta^* = 0$, and by (2.11) we must have $\varepsilon^* = 0$; therefore $F(x^*) = 0$ by (2.14). Now (2.10) and (2.8) yield

$$\frac{q}{\varepsilon_k^2} \Delta_k^2 + \|w\|^2 + 2(1 - q\Delta_k)^2 \sigma_k \beta_1 \leq \frac{1}{\varepsilon_k^2} \|F(x^k)\|^2.$$

Since $\beta_1 > 0$, the last term on the left-hand side tends to ∞ as $k \rightarrow \infty$. Thus the vectors $y^k := \varepsilon_k^{-1} F(x^k)$ satisfy $\|y^k\| \rightarrow \infty$, the vectors $z^k := y^k / \|y^k\|$ have norm 1, and (2.12) implies that the numbers μ_i^k ($i = 1, \dots, n$), defined by

$$\mu_i^k := \frac{1}{\|y^k\|} f_{x_i}(x^k) + \frac{1}{(1 - q\Delta_k)^2} (F'(x^k)^T z^k)_i - \frac{1}{(1 - q\Delta_k)^2 \|y^k\|} (F'(x^k)^T w)_i,$$

satisfy

$$\mu_i^k \begin{cases} \geq 0 & \text{if } x_i^k = u_i, \\ = 0 & \text{if } u_i < x_i^k < v_i, \\ \leq 0 & \text{if } x_i^k = v_i. \end{cases}$$

If we pick a convergent subsequence z^{k_l} with limit z^* and pass to the limit we obtain

$$(F'(x^*)^T z^*)_i \begin{cases} \geq 0 & \text{if } x_i^* = u_i, \\ = 0 & \text{if } u_i < x_i^* < v_i, \\ \leq 0 & \text{if } x_i^* = v_i. \end{cases}$$

Now similarly as above this yields $z^* = 0$, which is a contradiction to $\|z^*\| = 1$. Thus such a sequence $(x^k, \varepsilon_k, \sigma_k)$ cannot exist, and for sufficiently large σ all Kuhn–Tucker points of f_σ are of the form $(x, 0)$.

Now let (x^*, ε^*) be a local minimizer of f_σ with finite $f_\sigma(x^*, \varepsilon^*)$. If $\varepsilon^* > 0$ then (x^*, ε^*) must be a Kuhn–Tucker point, contradiction. Therefore, $\varepsilon^* = 0$, and since $f_\sigma(x^*, \varepsilon^*)$ is finite, $\Delta(x^*, \varepsilon^*) = 0$. Now (2.4) implies $F(x^*) = 0$, so that x^* is a feasible point of (2.1). Thus (2.6) implies that there is a neighborhood of x^* where $f(x) \geq f(x^*)$ for feasible x . Therefore x^* is a local minimizer of (2.1). \square

We conclude that under the stated assumptions, minimizing the penalty function f_σ for sufficiently large σ yields a minimizer of the original problem. Conversely, as we shall prove in §5 in a more general setting, a minimizer x^* of (2.1) yields a minimizer $(x^*, 0)$ of f_σ for sufficiently large σ and slightly stronger conditions on $\beta(\varepsilon)$.

Remark. If $w_i \neq 0$ for $i = 1, \dots, n$, we can write $w_i = \lambda_i^{-1}$ and rewrite (2.2) as

$$\begin{aligned} \min \quad & f(x) \\ \text{s.t.} \quad & \lambda_i F_i(x) = \varepsilon \quad (i = 1, \dots, m), \\ & x \in [u, v], \quad \varepsilon = 0. \end{aligned}$$

This is again of the form (2.2) with F_i replaced by $\lambda_i F_i$ and w_i replaced by 1. Therefore, the theorem also holds with $\Delta(x, \varepsilon) = \sum_{i=1}^m (\varepsilon - \lambda_i F_i(x, \varepsilon))^2$ in the penalty function

(2.3). Now the penalty function has an augmented Lagrangian interpretation. Indeed, for (x, ε) with small $\Delta(x, \varepsilon)$ we obtain

$$\begin{aligned} f_\sigma(x) &= f(x) + \frac{1}{2\varepsilon} \cdot \frac{\Delta(x, \varepsilon)}{1 - q\Delta(x, \varepsilon)} + \sigma\beta(\varepsilon) \\ &= f(x) + \frac{1}{2\varepsilon}\Delta(x, \varepsilon) + \sigma\beta(\varepsilon) + O(\Delta(x, \varepsilon)^2) \\ &= f(x) + \frac{1}{2\varepsilon} \sum_{i=1}^m (\lambda_i F_i(x) - \varepsilon)^2 + \sigma\beta(\varepsilon) + O(\Delta(x, \varepsilon)^2) \\ &= f(x) - \sum \lambda_i F_i(x) + \frac{1}{2\varepsilon} \sum \lambda_i^2 F_i(x)^2 + \frac{m\varepsilon}{2} + \sigma\beta(\varepsilon) + O(\Delta(x, \varepsilon)^2). \end{aligned}$$

Thus, for fixed ε and up to constant additive terms and higher order terms, f_σ is an augmented Lagrangian, and the λ_i play the role of (fixed, initial) Lagrange multiplier estimates.

In particular, as in traditional multiplier penalty functions [1], if the λ_i are close to the Lagrange multipliers at the optimizer x^* then $f_\sigma(x, \varepsilon)$ is nearly stationary at x^* for arbitrary fixed $\varepsilon > 0$. Therefore, numerical schemes for the minimization of $f_\sigma(x, \varepsilon)$ get close to the minimizer already when ε is not very small, and hence when the Hessian of f_σ (which gets more and more ill-conditioned as $\varepsilon \rightarrow 0$) is still well-conditioned.

3. Regularization of nonsmooth and ill-conditioned problems. In order to regularize the optimization problem (2.1) with not necessarily smooth functions f and F , we assume that we can embed f and F into a family of regularized functions $f(x, \varepsilon)$ and $F(x, \varepsilon)$ that are twice continuously differentiable in (x, ε) when $\varepsilon > 0$ and satisfy

$$f(x) = f(x, 0) = \lim_{\varepsilon \rightarrow 0} f(x, \varepsilon), \quad F(x) = F(x, 0) = \lim_{\varepsilon \rightarrow 0} F(x, \varepsilon).$$

Of course, the case where the objective function and the constraints are already well-behaved needs no modification, and in this case we simply put $f(x, \varepsilon) = f(x)$ and $F(x, \varepsilon) = F(x)$ for all ε .

Nonsmooth functions arising in practice are often *factorable*, i.e., composed of a finite sequence of elementary operations. Most elementary operations are smooth; the nonsmoothness arises through a small number of nonsmooth elementary functions. A natural regularization approach for factorable functions is to write each nonsmooth elementary function $N(x)$ as a limit of smooth functions $N(x, \varepsilon)$ that are twice continuously differentiable in (x, ε) when $\varepsilon > 0$,

$$N(x) = \lim_{\varepsilon \rightarrow 0} N(x, \varepsilon).$$

Assuming that the objective and constraint functions are factorable, we may replace each occurrence $N(r_i)$ of a nonsmooth elementary function in the definition of the objective and constraint functions with $N(r_i, \varepsilon\rho_i)$ depending on an intermediate result r_i and a suitable scaling constant ρ_i . Then we end up with regularized functions $f(x, \varepsilon)$ and $F(x, \varepsilon)$ with the required properties. Possible forms of $N(x, \varepsilon)$ for the most important nonsmooth $N(x)$ are given in Table 3.1. Note that the first two formulas are independent of ξ ; the particular choice indicated is numerically stable and allows to restrict the sum to those terms where the exponent is $> \log macheps$, where *macheps* is the machine accuracy.

Some smooth nonlinear programs are very difficult to solve since the Hessian matrix of the Lagrangian is severely ill-conditioned everywhere. Often, the reason for this is that the objective function or some constraint contains subexpressions involving some huge or tiny constants. Such constants can be regularized, too, by adapting them according to the last two lines of Table 3.1.

TABLE 3.1
Some regularization recipes

$N(x)$	$N(x, \varepsilon)$	condition
$\max_k x_k$	$\xi + \varepsilon \log \sum \exp((x_k - \xi)/\varepsilon)$	$\xi = \max_k x_k$
$\min_k x_k$	$\xi - \varepsilon \log \sum \exp((\xi - x_k)/\varepsilon)$	$\xi = \min_k x_k$
x^t ($x \geq 0, t < 2$)	$(x + \varepsilon)^t$	$t \neq 0, 1$
$x^t \log x$ ($x \geq 0, t \leq 2$)	$x^t \log(x + \varepsilon)$	$t = 0, 1, 2$
$ x ^t$ ($t < 2$)	$(x + \varepsilon)^t \log(x + \varepsilon)$	$t \neq 0, 1, 2$
$ x ^t \log x $ ($0 < t \leq 2$)	$ x ^{t+k} / (x ^k + \varepsilon^k)$	$k = \lceil 2 - t \rceil$
c (huge constant)	$ x ^{t+k} \log x / (x ^k + \varepsilon^k)$	$k = 1 + \lfloor 2 - t \rfloor$
c (tiny constant)	$c/(1 + \varepsilon c)$	
	$c + \varepsilon \operatorname{sign} c$	

To approximate elementary functions with step discontinuities, such as

$$\operatorname{pos}(x) = \begin{cases} 1 & \text{if } x > 0, \\ 0 & \text{if } x \leq 0, \end{cases} \quad \operatorname{nneg}(x) = \begin{cases} 1 & \text{if } x \geq 0, \\ 0 & \text{if } x < 0, \end{cases}$$

$$\operatorname{sign}(x) = \begin{cases} 1 & \text{if } x > 0, \\ 0 & \text{if } x = 0, \\ -1 & \text{if } x < 0, \end{cases}$$

one may use the regularizations given in Table 3.2 (using $x_+ = \max(x, 0)$). However, no accompanying theory is available since the results presented in §5 require the Lipschitz continuity of the functions involved.

TABLE 3.2
Regularization of functions with step discontinuities

$N(x)$	$N(x, \varepsilon)$	condition
$\operatorname{pos}(x)$	$x_+^3 / (x_+^3 + \varepsilon^3)$	
$\operatorname{nneg}(x)$	$x_+^3 / (x_+^3 + \varepsilon^3)$	
$\operatorname{sign}(x)$	$x^3 / (x ^3 + \varepsilon^3)$	
$\begin{cases} p & \text{if } x \geq 0 (> 0), \\ q & \text{otherwise} \end{cases}$	$\begin{cases} px^3 / (x^3 + \varepsilon^3) & \text{if } x \geq 0, \\ qx^3 / (x^3 - \varepsilon^3) & \text{otherwise} \end{cases}$	$pq \leq 0$

4. Regular zeros of nonsmooth functions. In this section we derive some technical results that are needed for the successful analysis of nonsmooth equality constraints.

DEFINITION 4.1. A point $x^* \in \mathbb{R}^n$ is called a regular zero of a function $H : D \subseteq \mathbb{R}^n \rightarrow \mathbb{R}^m$, $m \leq n$, if x^* is in the interior of D , satisfies $H(x^*) = 0$, and there are a closed, convex and bounded set \mathcal{A} of $m \times n$ -matrices and a matrix $B \in \mathbb{R}^{(n-m) \times n}$

such that the augmented matrix $\begin{pmatrix} A \\ B \end{pmatrix}$ is nonsingular for all $A \in \mathcal{A}$ and for every x, y in some neighborhood $N \subseteq D$ of x^* there exists a matrix $A \in \mathcal{A}$ with

$$(4.1) \quad H(x) - H(y) = A(x - y).$$

The regularity of a zero can be verified under quite general circumstances.

PROPOSITION 4.2. *A point $x^* \in \mathbb{R}^n$ is a regular zero of H if H is continuously differentiable in a neighborhood of x^* and $H'(x^*)$ has full rank.*

Proof. If $H'(x^*)$ has full rank, there is a matrix B such that $\begin{pmatrix} H'(x^*) \\ B \end{pmatrix}$ is square and nonsingular. By continuity, there exists a convex neighborhood N of x^* such that $\begin{pmatrix} H'(x) \\ B \end{pmatrix}$ is nonsingular for $x \in N$ and, as in the proof of [12, Proposition 5.1.4], (4.1) is satisfied if we take for \mathcal{A} the closed convex hull of $\{H'(x) \mid x \in N\}$. \square

The preceding result generalizes to certain piecewise differentiable functions if the nonsmoothness is not too severe. For example, we have

PROPOSITION 4.3. *A point $x^* \in \mathbb{R}^n$ is a regular zero of $H(x) = G(x, |x - x^*|)$ if G is continuously differentiable in a neighborhood of $(x^*, 0)$ with $G(x^*, 0) = 0$, and if there exists a matrix B such that, for all diagonal matrices $\Sigma \in \mathbb{R}^{n \times n}$ with $|\Sigma_{ii}| \leq 1$ for $i = 1, \dots, n$, the matrix $\begin{pmatrix} \partial_1 G(x^*, 0) + \partial_2 G(x^*, 0)\Sigma \\ B \end{pmatrix}$ is nonsingular.*

Proof. We mimic the proof of Neumaier [12, Proposition 5.1.4]. Without loss of generality, $x^* = 0$. There is a ball $N = B[0; \delta]$ such that $G(x, z)$ is continuously differentiable for $(x, z) \in N$ and every matrix $\begin{pmatrix} \partial_1 G(x, z) + \partial_2 G(x, z)\Sigma \\ B \end{pmatrix}$ is nonsingular. Let $x, y \in N$. By a version of the mean value theorem we have

$$\begin{aligned} G(x, |x|) - G(y, |y|) &= \int_0^1 \partial_1 G(y + s(x - y), |x|)(x - y) ds \\ &\quad + \int_0^1 \partial_2 G(y, |y| + s(|x| - |y|))(|x| - |y|) ds \\ &= A(x - y), \end{aligned}$$

where

$$A = \int_0^1 \partial_1 G(y + s(x - y), |x|) ds + \int_0^1 \partial_2 G(y, |y| + s(|x| - |y|))\Sigma ds$$

and Σ is the diagonal matrix with the diagonal entries

$$\Sigma_{ii} := \begin{cases} (|x_i| - |y_i|)/(x_i - y_i) & \text{if } x_i \neq y_i, \\ 0 & \text{otherwise.} \end{cases}$$

Since N is convex, A is contained in the closed convex hull \mathcal{A} of the set of expressions $\partial_1 G(x, |x'|) + \partial_2 G(y, |y'|)\Sigma$, where $x, x', y, y' \in N$ and Σ is a diagonal matrix with $|\Sigma_{ii}| \leq 1$ for $i = 1, \dots, n$. If N is chosen sufficiently small, $\begin{pmatrix} A \\ B \end{pmatrix}$ is nonsingular for all $A \in \mathcal{A}$. \square

In particular, this applies to $H(x) = G(x) + \delta G_1(x, |x|)$ if G and G_1 are continuously differentiable near 0, $G'(0)$ has full rank, and δ is sufficiently small.

With a similar argument but now involving generalized derivatives (essentially the convex hull of the limit set of gradients of nicely approximating smooth functions; see, e.g., [2, 12]), the following result can be proved.

PROPOSITION 4.4. *A point $x^* \in \mathbb{R}^n$ is a regular zero of $H(x)$ if H is Lipschitz continuous in a neighborhood of x^* with $H(x^*) = 0$, and if there exists a matrix B such that, for all matrices H' contained in the generalized derivative of H at x^* , the matrix $\begin{pmatrix} H' \\ B \end{pmatrix}$ is nonsingular.*

For x near a regular zero of H and an arbitrary set J of indices, one can find a small perturbation of the order of $O(\|H_J(x)\|)$ such that the perturbed vector y satisfies $H_J(y) = 0$ and $H_i(y) = H_i(x)$ for all $i \notin J$.

THEOREM 4.5. *Let x^* be a regular zero of $H : D \subseteq \mathbb{R}^n \rightarrow \mathbb{R}^m$, $m \leq n$. Then there are a neighborhood $N_0 \subseteq N$ of x^* and a constant $\gamma_0 > 0$ such that for each $x \in N_0$ and each subset J of $\{1, \dots, m\}$ there exists a vector $y = y(x) \in N$ with $H_i(y) = 0$ for $i \in J$ and $H_i(y) = H_i(x)$ for $i \notin J$ such that*

$$\|x - y\| \leq \gamma_0 \|H_J(x)\|.$$

Proof. We define the neighborhood $N_0 := B[x^*; (2\gamma_0(L + \|B\|))^{-1}r] \cap N$, where $r > 0$ is such that the closed ball $N_1 := B[x^*; r]$ is contained in the neighborhood N of x^* , and

$$L := \sup_{A \in \mathcal{A}} \|A\| < \infty, \quad \gamma_0 := \sup_{A \in \mathcal{A}} \left\| \begin{pmatrix} A \\ B \end{pmatrix}^{-1} \right\| < \infty$$

by assumption. By (4.1), the constant L is a Lipschitz constant for H . We fix a vector $x \in N_0$ and a subset J of $\{1, \dots, m\}$, and put $K := \{1, \dots, m\} \setminus J$.

To find y , we want to apply the nonsmooth inverse function theorem given in Neumaier [12, Theorem 5.1.6(iv)] to the mapping $F : N \rightarrow \mathbb{R}^n$ defined by

$$F(z) := \begin{pmatrix} H(z) \\ B(z - x^*) \end{pmatrix} \quad \text{for } z \in N,$$

with the right-hand side

$$(4.2) \quad b^* = \begin{pmatrix} b \\ B(x - x^*) \end{pmatrix},$$

where $b_i = 0$ for $i \in J$ and $b_i = H_i(x)$ for $i \in K$, and $\tilde{x}^0 = x^*$. Since $H(x^*) = 0$ implies $F(x^*) = 0$, this requires that we show

$$(4.3) \quad F(z) \neq sb^* \quad \text{for all } z \in \partial N_1, s \in [0, 1).$$

If this holds, [12, Theorem 5.1.6(iv)] implies that there is a unique $y \in N_1$ with $F(y) = b^*$, hence $H_i(y) = 0$ for $i \in J$ and $H_i(y) = H_i(x)$ for $i \in K$. Moreover,

$$F(x) - F(y) = \begin{pmatrix} A \\ B \end{pmatrix} (x - y)$$

for some $A \in \mathcal{A}$ and

$$\|F(x) - F(y)\| = \|H_J(x)\|,$$

hence

$$\|x - y\| \leq \left\| \begin{pmatrix} A \\ B \end{pmatrix}^{-1} \right\| \|H_J(x)\| \leq \gamma_0 \|H_J(x)\|.$$

Thus y has the required properties.

To show (4.3), we suppose that $F(z) = sb^*$ for some $z \in N_1$, $s \in [0, 1)$. Since N_1 is contained in N , there exists an $A \in \mathcal{A}$ such that

$$sb^* = F(z) = F(z) - F(x^*) = \begin{pmatrix} A \\ B \end{pmatrix} (z - x^*),$$

and we obtain

$$\|z - x^*\| = \|s \begin{pmatrix} A \\ B \end{pmatrix}^{-1} b^*\| \leq s\gamma_0 \|b^*\|.$$

Since $H(x^*) = 0$, (4.2) implies $\|b^*\| \leq \|H_K(x) - H_K(x^*)\| + \|B(x - x^*)\| \leq (L + \|B\|)\|x - x^*\|$, and we conclude

$$(4.4) \quad \|z - x^*\| \leq s\gamma_0(L + \|B\|)\|x - x^*\|.$$

Hence $z \in B[x^*, r/2]$ cannot lie on the boundary of N_1 ; in particular, (4.3) holds. This proves the theorem. \square

5. The local exactness proof. We now consider the optimization problem (2.1), where we now allow f and F to be nonsmooth functions. Clearly, with the embedding of §3 and a fixed $w \in \mathbb{R}^m$, (2.1) is equivalent to

$$(5.1) \quad \begin{aligned} \min \quad & f(x, \varepsilon) \\ \text{s.t.} \quad & F_i(x, \varepsilon) = \varepsilon w_i \quad (i = 1, \dots, m), \\ & x \in [u, v], \quad \varepsilon = 0. \end{aligned}$$

We assume that f and F are continuous on $D \times [0, \bar{\varepsilon}]$, where D is an open set containing $[u, v]$ and $\bar{\varepsilon} > 0$, and twice continuously differentiable on its interior.

We use again the expression (2.3) for the penalty function but with $f(x, \varepsilon)$ in place of $f(x)$ and the regularized constraint violation measure

$$(5.2) \quad \Delta(x, \varepsilon) := \|\varepsilon w - F(x, \varepsilon)\|^2.$$

If, in addition to the assumptions mentioned after (2.4), β is twice continuously differentiable for $\varepsilon > 0$, the function f_σ is twice continuously differentiable for $(x, \varepsilon) \in (0, \bar{\varepsilon}) \times [u, v]$ with $\Delta(x, \varepsilon) < q^{-1}$.

It is conceivable that for this penalty function, a suitable analogue of Theorem 2.1 holds even in the nonsmooth case, but this requires an extension of the Kuhn–Tucker optimality conditions to nonsmooth problems, and we have not tried to handle the technicalities associated with this. On the other hand, we show here (after some preparation) that a converse of Theorem 2.1 can be proved in the nonsmooth case.

Assumptions. For the formal analysis, we shall suppose that, in addition to the general assumptions made above, the following assumptions (H_f) , (H_F) and (H_ε) are satisfied in a neighborhood of some local (or global) minimizer $x = x^*$ of (5.1). Let $I := \{i \mid u_i < x_i^* < v_i\}$; to simplify notation we assume that $I = \{1, \dots, p\}$ with $m \leq p \leq n$.

(H_f) $f(\cdot, 0)$ is Lipschitz continuous with the Lipschitz constant k .

(H_F) x^* is a regular zero of $F(\cdot, 0)$ and x_I^* is a regular zero of $G : D_1 \subseteq \mathbb{R}^p \rightarrow \mathbb{R}^m$, defined by $G(\tilde{x}) := F(x, 0)$, where $x_i = \tilde{x}_i$ for $i = 1, \dots, p$ and $x_i = x_i^*$ for $i = p + 1, \dots, n$ and D_1 is an appropriate open set containing $[u_I, v_I]$. N is a neighborhood of x^* according to Definition 4.1 such that, in addition, $F(\cdot, 0)$ is Lipschitz continuous in N and $x_I \in [u_I, v_I]$ for $x \in N$.

(H_ε) There are positive constants $\bar{\varepsilon}$ and K such that for all $x \in N$ and all $\varepsilon \in [0, \bar{\varepsilon}]$,

$$\|F(x, 0) - F(x, \varepsilon)\|_\infty \leq K\varepsilon, \quad |f(x, 0) - f(x, \varepsilon)| \leq K\varepsilon.$$

(H_β) β satisfies $\liminf_{\varepsilon \rightarrow 0} \beta(\varepsilon)/\sqrt{\varepsilon} > 0$.

LEMMA 5.1. x^* is a regular zero of $H : D \subseteq \mathbb{R}^n \rightarrow \mathbb{R}^{m+n-p}$, defined by $H_i(x) := F_i(x, 0)$ for $i = 1, \dots, m$ and $H_i(x) := x_{i-m+p} - x_{i-m+p}^*$ for $i = m+1, \dots, m+n-p$, if and only if x^* is a regular zero of $F(\cdot, 0)$ and x_I^* is a regular zero of the mapping G defined in (H_F).

Proof. Let $x, y \in N$, where N is an appropriate neighborhood of x^* according to Definition 4.1. In both cases we have $F(x, 0) - F(y, 0) = A(x - y)$ for a matrix $A \in \mathcal{A}$, where \mathcal{A} is a closed, convex and bounded set of $m \times n$ -matrices such that the augmented matrix $\begin{pmatrix} A \\ B \end{pmatrix}$ is nonsingular for all $A \in \mathcal{A}$ for some matrix $B \in \mathbb{R}^{(n-m) \times n}$. Then

$$H(x) - H(y) = \begin{pmatrix} A \\ A' \end{pmatrix} (x - y),$$

where $A' := (0 \ I_{n-p}) \in \mathbb{R}^{(n-p) \times n}$, and

$$G(x_I) - G(y_I) = A_{:,I}(x_I - y_I).$$

Let x^* be a regular zero of H and let $B' \in \mathbb{R}^{(p-m) \times n}$ be such that $\begin{pmatrix} A \\ A' \\ B' \end{pmatrix}$ is nonsingular for all $A \in \mathcal{A}$. Then $\begin{pmatrix} A_{:,I} \\ B'_{:,I} \end{pmatrix}$ is nonsingular, i.e., x_I^* is a regular zero of G . Conversely, let x_I^* be a regular zero of G and let $B'_{:,I}$ be such that $\begin{pmatrix} A_{:,I} \\ B'_{:,I} \end{pmatrix}$ is nonsingular. Then $\begin{pmatrix} A \\ A' \\ B'' \end{pmatrix}$ is nonsingular with $B'' = (B'_{:,I} \ 0) \in \mathbb{R}^{(p-m) \times n}$. Therefore x^* is a regular zero of H . \square

LEMMA 5.2. If (H_f) and (H_F) hold, there are a neighborhood $N_0 \subseteq N$ of x^* and a constant $\gamma_1 > 0$ such that

$$f(x, 0) \geq f(x^*, 0) - \gamma_1 \|F(x, 0)\| \quad \text{for all } x \in N_0 \cap [u, v].$$

Proof. By (H_F), x^* is a regular zero of the mapping H defined in Lemma 5.1. Let $N_0 \subseteq N$ and γ_0 be as in Theorem 4.5, and let $x \in N_0$. Then by Theorem 4.5 with $J := \{i \mid F_i(x, 0) \neq 0\}$ there exists a $y = y(x)$ with $H(y) = 0$ such that

$$\|x - y\| \leq \gamma_0 \|H_J(x)\| = \gamma_0 \|F(x, 0)\|.$$

The fact that $y_i = x_i^*$ for $i = p + 1, \dots, n$ and the choice of N in (H_F) imply that $y \in [u, v]$, i.e., y is a feasible point. We therefore have $f(y, 0) \geq f(x^*, 0)$ by assumption, and

$$\begin{aligned} f(x, 0) &= f(x, 0) - f(y, 0) + f(y, 0) \geq f(x^*, 0) - k\|x - y\| \\ &\geq f(x^*, 0) - k\gamma_0\|F(x, 0)\|, \end{aligned}$$

which finishes the proof. \square

Now we can prove the main theorem of this section.

THEOREM 5.3. *Under the above assumptions and for sufficiently large σ , there are a neighborhood N' of x^* and an $\varepsilon' \in (0, \bar{\varepsilon}]$ such that*

$$f_\sigma(x, \varepsilon) > f_\sigma(x^*, 0) = f(x^*, 0) \quad \text{for all } (x, \varepsilon) \in N' \times (0, \varepsilon'].$$

In particular, $(x^*, 0)$ is a local minimizer of f_σ .

Proof. Let the neighborhood $N' \subseteq N_0$ of x^* (N_0 as in Lemma 5.2) be sufficiently small such that

$$(5.3) \quad \sup_{x \in N'} (f(x^*, 0) - f(x, 0)) \leq \frac{1}{2},$$

let $\varepsilon' \in (0, \bar{\varepsilon}]$, $\varepsilon' \leq 1$ be sufficiently small such that

$$(5.4) \quad \beta(\varepsilon) \geq \beta_2\sqrt{\varepsilon}$$

for $0 \leq \varepsilon \leq \varepsilon'$ and a $\beta_2 > 0$, and let

$$(5.5) \quad \sigma \geq \frac{1}{\beta_2}(K(\gamma_1 + 1) + \gamma_1(\|w\| + 1)).$$

For $(x, \varepsilon) \in N' \times (0, \varepsilon']$ we distinguish two cases.

Case 1. Let $\Delta(x, \varepsilon) \geq \varepsilon$. Then

$$\begin{aligned} f_\sigma(x, \varepsilon) &\geq f(x, \varepsilon) + \frac{1}{2} + \sigma\beta(\varepsilon) \\ &\geq f(x^*, 0) - \frac{1}{2} - K\varepsilon + \frac{1}{2} + \sigma\beta(\varepsilon) \\ &\geq f(x^*, 0) - K\sqrt{\varepsilon} + \sigma\beta_2\sqrt{\varepsilon} > f(x^*, 0), \end{aligned}$$

where we have used (H_ε) , (5.3), (5.4), (5.5) and the fact that $\varepsilon \leq \varepsilon' \leq 1$.

Case 2. If $\Delta(x, \varepsilon) < \varepsilon$, then $\|F(x, \varepsilon)\| < \varepsilon\|w\| + \|\varepsilon w - F(x, \varepsilon)\| \leq \varepsilon\|w\| + \sqrt{\varepsilon}$, hence by Lemma 5.2 and (H_ε) ,

$$\begin{aligned} f(x^*, 0) &\leq f(x, 0) + \gamma_1\|F(x, 0)\| \\ &\leq f(x, \varepsilon) + K\varepsilon + \gamma_1(\|F(x, \varepsilon)\| + K\varepsilon) \\ &< f(x, \varepsilon) + K(\gamma_1 + 1)\varepsilon + \gamma_1(\sqrt{\varepsilon} + \varepsilon\|w\|). \end{aligned}$$

Therefore $f_\sigma(x, \varepsilon) \geq f(x, \varepsilon) + \sigma\beta(\varepsilon) > f(x^*, 0)$ by (5.4), (5.5) and $\varepsilon \leq \varepsilon' \leq 1$. \square

6. An example. To illustrate the theory developed, we consider the simple smooth nonlinear optimization problem

$$\begin{aligned} \min \quad & x_1^3 x_2^3 \\ \text{s.t.} \quad & x_1^2 + x_2^2 = 1. \end{aligned}$$

It has a bounded feasible domain, two global minimizers at $x^* = (\frac{1}{2}\sqrt{2}, -\frac{1}{2}\sqrt{2})^T$ and $x^{**} = (-\frac{1}{2}\sqrt{2}, \frac{1}{2}\sqrt{2})^T$ with $f(x^*) = f(x^{**}) = -\frac{1}{8}$, and no other local minima. The traditional quadratic penalty function for this problem,

$$p(x) = x_1^3 x_2^3 + \frac{1}{2\varepsilon}(x_1^2 + x_2^2 - 1)^2,$$

is unbounded below for all $\varepsilon > 0$ since, e.g., $p(x) \rightarrow -\infty$ for $x = (s, -s)^T$, $s \rightarrow \infty$. This is also the case for traditional exact penalty functions, including multiplier penalty functions [1] that use an additional term $+\lambda(x_1^2 + x_2^2 - 1)$. On the other hand, our new penalty function is bounded below. For $w = 1$ it reads

$$f_\sigma(x, \varepsilon) = \begin{cases} x_1^3 x_2^3 & \text{for } \varepsilon = \Delta(x, \varepsilon) = 0, \\ x_1^3 x_2^3 + \frac{1}{2\varepsilon} \cdot \frac{r^2}{1 - qr^2} + \sigma\beta(\varepsilon) & \text{for } \varepsilon > 0, |r| < q^{-1/2}, \\ \infty & \text{otherwise,} \end{cases}$$

where $r := 1 + \varepsilon - x_1^2 - x_2^2$. Since $f_\sigma(x, \varepsilon) = \infty$ if $\|x\| \geq \sqrt{q^{-1/2} + 1 + \varepsilon}$, the boundedness of our penalty function below is trivial. The Mangasarian–Fromovitz condition holds for all $x \neq 0$, hence the assumptions of Theorem 2.1 are satisfied if $q^{-1/2} + \bar{\varepsilon} < 1$ and (2.8) holds; in this case, every local minimizer of the penalty function with a sufficiently large σ gives a solution of the original constrained problem.

In this particular example we can give an explicit analysis, and find that in fact weaker assumptions than those given in Theorem 2.1 suffice to get the conclusions, since $\bar{\varepsilon}$ can be chosen arbitrarily large. To show this, let (x, ε) be a Kuhn–Tucker point of f_σ with $\varepsilon > 0$ and $|r| < q^{-1/2}$. Then $\partial f_\sigma / \partial x_1$ and $\partial f_\sigma / \partial x_2$ vanish at (x, ε) , and $\partial f_\sigma(x, \varepsilon) / \partial \varepsilon \leq 0$, with equality if $\varepsilon < \bar{\varepsilon}$. We have

$$\begin{aligned} \frac{\partial f_\sigma}{\partial x_1}(x, \varepsilon) &= x_1 \left(3x_1 x_2^3 - \frac{2r}{\varepsilon(1 - qr^2)^2} \right), \\ \frac{\partial f_\sigma}{\partial x_2}(x, \varepsilon) &= x_2 \left(3x_1^3 x_2 - \frac{2r}{\varepsilon(1 - qr^2)^2} \right), \\ \frac{\partial f_\sigma}{\partial \varepsilon}(x, \varepsilon) &= \frac{r(2\varepsilon - r + qr^3)}{2\varepsilon^2(1 - qr^2)^2} + \sigma\beta'(\varepsilon). \end{aligned}$$

Since $\sigma\beta'(\varepsilon) > 0$, $\partial f_\sigma / \partial \varepsilon \leq 0$ implies

$$(6.1) \quad r(2\varepsilon - r + qr^3) < 0;$$

in particular $r \neq 0$. Under this condition, the other partial derivatives vanish iff either $x_1 = x_2 = 0$ or

$$(6.2) \quad 3x_1 x_2^3 = 3x_1^3 x_2 = \frac{2r}{\varepsilon(1 - qr^2)^2}.$$

If (6.2) holds then $x_1 = \pm x_2$, the definition of r gives $1 + \varepsilon - r = 2x_1^2$, and (6.2) simplifies to

$$(6.3) \quad 8|r| = 3\varepsilon(1 - qr^2)^2(1 + \varepsilon - r)^2.$$

If $r > 0$ then $r > 2\varepsilon + qr^3 > 2\varepsilon$ by (6.1), giving $0 \leq 1 + \varepsilon - r < 1 - \varepsilon < 1$. Now (6.3) implies $8r \leq 3\varepsilon$, contradicting $r > 2\varepsilon$. And if $r < 0$ then (6.3) and $qr^2 < 1$ imply for $q \geq 1$ that

$$\frac{-r}{2\varepsilon(1 - qr^2)^2} = \frac{3}{16}(1 + \varepsilon + |r|)^2 \leq \frac{3}{16}(2 + \bar{\varepsilon}) =: t$$

and

$$2\varepsilon - r + qr^3 \leq 2\varepsilon - r \leq (2 + 2t)\varepsilon.$$

Now $\partial f_\sigma / \partial \varepsilon \leq 0$ gives $\sigma\beta'(\varepsilon) \leq t(2 + 2t)$. If (2.8) holds and $\sigma > 2t(t + 1)/\beta_1$, this is violated, (6.2) is impossible, and the only Kuhn-Tucker point of f_σ with $\varepsilon > 0$ can be at $x_1 = x_2 = 0$. But then $r = 1 + \varepsilon$ and (6.1) requires $q < (1 - \varepsilon)/(1 + \varepsilon)^3$, again impossible if $q \geq 1$.

Thus, for this example, the conclusion of Theorem 2.1 is valid for arbitrary $\bar{\varepsilon} > 0$ if (2.8) holds and $q \geq 1$, provided that σ is sufficiently large. Note that for $q \geq 1$ we have $\Delta(0, \varepsilon) = (1 + \varepsilon)^2 > q^{-1}$, so that the only point violating the Mangasarian–Fromovitz condition does not satisfy $\Delta(x, \varepsilon) < q^{-1}$.

We now look at the converse. Theorem 5.3 gives conditions guaranteeing that every solution of the constrained problem is a local minimizer of f_σ . To verify this explicitly we need to investigate when it is possible that f_σ is below the common value $-\frac{1}{8}$ of $f_\sigma(x^*, 0)$ and $f_\sigma(x^{**}, 0)$. Thus suppose that $f_\sigma(x, \varepsilon) \leq -\frac{1}{8}$, $(x, \varepsilon) \neq (x^*, 0), (x^{**}, 0)$. Then $\varepsilon > 0$. Since

$$x_1^3 x_2^3 \geq -\frac{1}{8}(x_1^2 + x_2^2)^3 = -\frac{1}{8}(1 + \varepsilon - r)^3 \geq -\frac{1}{8}(1 + \varepsilon + |r|)^3,$$

we find

$$(6.4) \quad 8\sigma\beta(\varepsilon) \leq -1 + (1 + \varepsilon + |r|)^3 - 4r^2/\varepsilon =: p(|r|), \quad |r| < q^{-1/2}.$$

Now $p(r)$ is a cubic polynomial. If $\varepsilon < \varepsilon_0 := \frac{1}{6}(-3 + \sqrt{33}) = 0.45742\dots$, the positive solution of $\varepsilon_0(1 + \varepsilon_0) = \frac{2}{3}$, then $\delta := 2 - 3\varepsilon(1 + \varepsilon) > 0$, and $p(r)$ has a unique local maximum at

$$r_0 = \frac{3\varepsilon(1 + \varepsilon)^2}{2 + \delta + \sqrt{8\delta}} \leq \frac{3}{2}\varepsilon(1 + \varepsilon)^2 \leq \frac{3}{2}\varepsilon(1 + \varepsilon_0)^2 = \varepsilon(1 + \varepsilon_0^{-1})$$

with function value

$$p(r_0) \leq -1 + (1 + \varepsilon + r_0)^3 \leq -1 + (1 + \varepsilon(2 + \varepsilon_0^{-1}))^3 \leq \frac{-1 + (2 + 2\varepsilon_0)^3}{\varepsilon_0} \varepsilon < 52\varepsilon.$$

Here we used the fact that $(-1 + (1 + \varepsilon(2 + \varepsilon_0^{-1}))^3)/\varepsilon$ is monotone increasing. Now

$$p(|r|) \leq \max(p(r_0), p(q^{-1/2})) < 52\varepsilon \quad \text{for } |r| < q^{-1/2}$$

if $p(q^{-1/2}) < 0$, which holds for any fixed q if ε is small enough. Thus (6.4) is violated if $\sigma\beta(\varepsilon) > 6.5\varepsilon$ and ε is small enough. If (H_β) holds, this is the case for arbitrary σ and sufficiently small $\varepsilon > 0$. Thus we have a contradiction, and the conclusion of Theorem 5.3 holds for arbitrary σ . We also see that in this example the conclusion of Theorem 5.3 still holds if the condition $\liminf_{\varepsilon \rightarrow 0} \beta(\varepsilon)/\sqrt{\varepsilon} > 0$ in (H_β) is relaxed to $\liminf_{\varepsilon \rightarrow 0} \beta(\varepsilon)/\varepsilon > 0$ and σ is chosen sufficiently large.

7. More general constraints. In this section we extend our theory to the more general constrained optimization problem

$$(7.1) \quad \begin{array}{ll} \min & f(x) \\ \text{s.t.} & x \in [u, v], \quad F_l \leq F(x) \leq F_u, \end{array}$$

where $F_l \in (\mathbb{R} \cup \{-\infty\})^m$ and $F_u \in (\mathbb{R} \cup \{\infty\})^m$ are vectors containing proper or infinite bounds on the constraint functions and $F_l \leq F_u$. This formulation is quite general since equality constraints are allowed by taking equal lower and upper bounds, and one-sided inequalities by taking infinite lower or upper bounds. Thus we treat equality constraints and one- or two-sided inequality constraints on the same footing. Moreover, we again assume that f and F are embedded into families of regularized functions $f(x, \varepsilon)$ and $F(x, \varepsilon)$ with $f(x, 0) = f(x)$ and $F(x, 0) = F(x)$.

For the one-dimensional boxes (which are just the closed intervals) we define the *magnitude* [12]

$$\langle \mathbf{x} \rangle := \min\{|x| \mid x \in \mathbf{x}\} = \begin{cases} \underline{x} & \text{if } \underline{x} > 0, \\ -\bar{x} & \text{if } \bar{x} < 0, \\ 0 & \text{otherwise.} \end{cases}$$

We also need the simple interval operations

$$\alpha + \beta \mathbf{x} = \{\alpha + \beta x \mid x \in \mathbf{x}\} = \begin{cases} [\alpha + \beta \underline{x}, \alpha + \beta \bar{x}] & \text{if } \beta \geq 0, \\ [\alpha + \beta \bar{x}, \alpha + \beta \underline{x}] & \text{if } \beta \leq 0. \end{cases}$$

Using this interval notation, we introduce a box-valued function \mathbf{E} on $[u, v] \times [0, \bar{\varepsilon}]$ by

$$(7.2) \quad \mathbf{E} : (x, \varepsilon) \rightarrow \mathbf{E}(x, \varepsilon) := F(x, \varepsilon) - [F_l, F_u],$$

where we assume that (H_ε) is satisfied, and consider the following optimization problem in inclusion form,

$$(7.3) \quad \begin{array}{ll} \min & f(x, \varepsilon) \\ \text{s.t.} & \varepsilon w_i \in \mathbf{E}_i(x, \varepsilon) \quad (i = 1, \dots, m), \\ & x \in [u, v], \quad \varepsilon = 0, \end{array}$$

where again $w \in \mathbb{R}^m$ is fixed. Clearly, this formulation is equivalent to the optimization problem (7.1). The usefulness of this particular formulation will become apparent when we look at the associated penalty function.

The penalty function is again defined by (2.3) with $f(x, \varepsilon)$ in place of $f(x)$, but now the constraint violation measure $\Delta(x, \varepsilon)$ is of the form

$$(7.4) \quad \Delta(x, \varepsilon) := \sum_{i=1}^m \Delta_i(x, \varepsilon)$$

with

$$(7.5) \quad \Delta_i(x, \varepsilon) := \langle \varepsilon w_i - \mathbf{E}_i(x, \varepsilon) \rangle^2,$$

the squared distance of εw_i from the interval $\mathbf{E}_i(x, \varepsilon)$. Without loss of generality we assume that the constraints are inequality constraints for $i = 1, \dots, r$ and equality constraints for $i = r + 1, \dots, m$, where $1 \leq r \leq m$. Then

$$\Delta_i(x, \varepsilon) = (\varepsilon w_i - (F_i(x, \varepsilon) - F_{li}))^2 \quad \text{for } i = r + 1, \dots, m$$

and

$$(7.6) \quad \Delta_i(x, \varepsilon) = (\varepsilon w_i - (F_i(x, \varepsilon) - y_i))^2 \quad \text{for } i = 1, \dots, r,$$

where

$$(7.7) \quad y_i = \begin{cases} F_i(x, \varepsilon) - \varepsilon w_i & \text{if } F_i(x, \varepsilon) - \varepsilon w_i \in [F_{li}, F_{ui}], \\ F_{li} & \text{if } F_i(x, \varepsilon) - \varepsilon w_i < F_{li}, \\ F_{ui} & \text{if } F_i(x, \varepsilon) - \varepsilon w_i > F_{ui}. \end{cases}$$

Moreover, $\Delta(x, \varepsilon)$ is continuously differentiable for $\varepsilon > 0$, with

$$\frac{\partial}{\partial x} \Delta_i(x, \varepsilon) = -2(\varepsilon w_i - F_i(x, \varepsilon) + y_i) \frac{\partial F_i}{\partial x}(x, \varepsilon)$$

and

$$\frac{\partial}{\partial \varepsilon} \Delta_i(x, \varepsilon) = 2(\varepsilon w_i - F_i(x, \varepsilon) + y_i)(w_i - \frac{\partial F_i}{\partial \varepsilon}(x, \varepsilon))$$

for $i = 1, \dots, r$.

We reduce this more general situation to the previous one with the aid of slack variables and use the abbreviations $J := \{1, \dots, r\}$ and $J' := \{r+1, \dots, m\}$. By introducing the slack variables $y_i := F_i(x, \varepsilon)$ for $i = 1, \dots, r$ we obtain the problem

$$(7.8) \quad \begin{aligned} \min & \quad f(x, \varepsilon) \\ \text{s.t.} & \quad x \in [u, v], \quad y \in [F_{lJ}, F_{uJ}], \quad \varepsilon = 0, \\ & \quad F_J(x, \varepsilon) - y = \varepsilon w_J, \quad F_{J'}(x, \varepsilon) - F_{lJ'} (= F_{J'}(x, \varepsilon) - F_{uJ'}) = \varepsilon w_{J'}, \end{aligned}$$

which is of the form (5.1). The penalty function for this problem is given by

$$\tilde{f}_\sigma(x, y, \varepsilon) = \begin{cases} f(x) & \text{for } \varepsilon = \tilde{\Delta}(x, y, \varepsilon) = 0, \\ f(x, \varepsilon) + \frac{1}{2\varepsilon} \cdot \frac{\tilde{\Delta}(x, y, \varepsilon)}{1 - q\tilde{\Delta}(x, y, \varepsilon)} + \sigma\beta(\varepsilon) & \text{for } \varepsilon > 0, \tilde{\Delta}(x, y, \varepsilon) < q^{-1}, \\ \infty & \text{otherwise,} \end{cases}$$

where

$$\tilde{\Delta}(x, y, \varepsilon) = \sum_{i=1}^r (\varepsilon w_i - (F_i(x, \varepsilon) - y_i))^2 + \sum_{i=r+1}^m (\varepsilon w_i - (F_i(x, \varepsilon) - F_{li}))^2.$$

Let $\tilde{E}_i(x, y) := F_i(x) - y_i$, $i = 1, \dots, r$, and $\tilde{E}_i(x, y) := F_i(x) - F_{li}$, $i = r+1, \dots, m$. Then

$$\tilde{E}'(x, y) = \begin{pmatrix} F'_J(x) & -I_r \\ F'_{J'}(x) & 0 \end{pmatrix},$$

and, for $m \leq n + r$, $\tilde{E}'(x, y)$ has full rank if and only if $F'_{J'}(x)$ has full rank.

LEMMA 7.1. *Let $(x, \varepsilon) \in [u, v] \times [0, \bar{\varepsilon}]$ with $\Delta(x, \varepsilon) < q^{-1}$. Then*

$$(7.9) \quad \Delta(x, \varepsilon) = \min_{y \in [F_{lJ}, F_{uJ}]} \tilde{\Delta}(x, y, \varepsilon)$$

and thus

$$f_\sigma(x, \varepsilon) = \min_{y \in [F_{lJ}, F_{uJ}]} \tilde{f}_\sigma(x, y, \varepsilon).$$

Moreover, if (x, ε) is a Kuhn–Tucker point of f_σ then there exists a vector $y \in [F_{lJ}, F_{uJ}]$ such that (x, y, ε) is a Kuhn–Tucker point of \tilde{f}_σ .

Proof. Let $(x, \varepsilon) \in [u, v] \times [0, \bar{\varepsilon}]$ with $\Delta(x, \varepsilon) < q^{-1}$, and let y be defined by (7.7). Then, for $i = r + 1, \dots, m$ we have $\Delta_i(x, \varepsilon) = (\varepsilon w_i - (F_i(x, \varepsilon) - F_{li}))^2$, and $\Delta_i(x, \varepsilon)$ is given by (7.6) and (7.7) for $i = 1, \dots, r$, i.e., $\Delta(x, \varepsilon) = \tilde{\Delta}(x, y, \varepsilon)$ for y defined by (7.7) and clearly (7.9) holds. Moreover, $\frac{\partial}{\partial x} \Delta(x, \varepsilon) = \frac{\partial}{\partial x} \tilde{\Delta}(x, y, \varepsilon)$ and $\frac{\partial}{\partial \varepsilon} \Delta(x, \varepsilon) = \tilde{\Delta}(x, y, \varepsilon)$. If (x, ε) is a Kuhn–Tucker point of f_σ , (x, y, ε) thus fulfils the Kuhn–Tucker conditions for x and ε , and due to the definition of y and the fact that

$$\frac{\partial \tilde{f}_\sigma}{\partial y_i}(x, y, \varepsilon) = \frac{1}{\varepsilon(1 - q\tilde{\Delta}(x, y, \varepsilon))^2}(\varepsilon w_i - F_i(x, \varepsilon) + y_i)$$

it also satisfies the Kuhn–Tucker conditions with respect to y . \square

The Mangasarian–Fromovitz condition for this problem holds for $(x, y) \in [u, v] \times [F_{lJ}, F_{uJ}]$ if

$$\begin{aligned} &F'_{J'}(x) \text{ has full rank, and there is a } p \in \mathbb{R}^n \text{ with } F'_{J'}(x)p = 0, \\ &p_i \begin{cases} > 0 & \text{if } x_i = u_i, \\ < 0 & \text{if } x_i = v_i, \end{cases} \\ &(F'_J(x)p)_i \begin{cases} > 0 & \text{if } y_i = F_{li}, \\ < 0 & \text{if } y_i = F_{ui}. \end{cases} \end{aligned}$$

Let $I := \{i \mid u_i < x_i^* < v_i\}$ and $J_1 := \{i \in J \mid F_i(x^*, 0) = F_{il} \text{ or } F_i(x^*, 0) = F_{iu}\}$; without loss of generality $I := \{1, \dots, p\}$. Moreover, let $E(x) := F(x, 0) - F(x^*, 0)$, $x \in D$. Then (H_F) is replaced by

(H_E) E is Lipschitz continuous in a neighborhood of x^* , x^* is a regular zero of $E_{J'}$: $D \subseteq \mathbb{R}^n \rightarrow \mathbb{R}^{m-r}$, and x_I^* is a regular zero of $G: D_1 \subseteq \mathbb{R}^p \rightarrow \mathbb{R}^{m-r+r_1}$, $r_1 := |J_1|$, defined by $G(\tilde{x}) := E_{J_1 \cup J'}(x)$ with $x_i := \tilde{x}_i$ for $i = 1, \dots, p$ and $x_i := x_i^*$ for $i = p + 1, \dots, n$, where D_1 is an appropriate open set containing $[u_I, v_I]$.

THEOREM 7.2. *If the above Mangasarian–Fromovitz condition holds for all $(x, y) \in [u, v] \times [F_{lJ}, F_{uJ}]$ with*

$$\sum_{i=1}^r (F_i(x) - y_i)^2 + \sum_{i=r+1}^m (F_i(x) - F_{li})^2 \leq (q^{-1/2} + \bar{\varepsilon}\|w\|)^2,$$

the set of these (x, y) is bounded and (2.8) is satisfied, the conclusions of Theorem 2.1 hold for the smooth case of the penalty function defined in this section. Moreover, under the assumptions (H_f) , (H_E) , (H_ε) and (H_β) the conclusions of Theorem 5.3 are satisfied for the penalty function defined by (2.3) with $f(x)$ replaced by $f(x, \varepsilon)$ and Δ defined by (7.4) and (7.5).

Acknowledgment. The authors gratefully acknowledge partial support by the Austrian Fonds zur Förderung der wissenschaftlichen Forschung (FWF) under grant P11516-MAT.

REFERENCES

- [1] D. P. BERTSEKAS, *Constrained Optimization and Lagrange Multiplier Methods*, Academic Press, New York, 1982.
- [2] F. H. CLARKE, *Optimization and Nonsmooth Analysis*, Wiley, New York, 1983.

- [3] A. R. CONN, N. I. M. GOULD, AND PH. L. TOINT, *Methods for nonlinear constraints in optimization calculations*, in The State of the Art in Numerical Analysis, I. S. Duff and G. A. Watson, eds., Clarendon Press, Oxford, 1997, pp. 363–390.
- [4] Y. M. ERMOLIEV, A. V. KRYAZHIMSKII, AND A. RUSZCZYŃSKI, *Constraint aggregation principle in convex optimization*, Math. Programming, 76 (1997), pp. 353–372.
- [5] K. KIWIEL, *Proximity control in bundle methods for convex nondifferentiable minimization*, Math. Programming, 46 (1990), pp. 105–122.
- [6] A. V. KUNTSEVICH AND F. KAPPEL, *SolvOpt – The Solver for Local Nonlinear Optimization Problems (Version 1.1, Matlab, C, FORTRAN)*, University of Graz, Graz, 1997, WWW document, <http://www.uni-graz.at/imawww/kuntsevich/solvopt/>.
- [7] C. LEMARÉCHAL, A. NEMIROVSKII, AND Y. NESTEROV, *New variants of bundle methods*, Math. Programming, 69 (1995), pp. 111–147.
- [8] C. LEMARÉCHAL AND C. SAGASTIZÁBAL, *Variable metric bundle methods: From conceptual to implementable forms*, Math. Programming, 76 (1997), pp. 393–410.
- [9] ———, *Practical aspects of the Moreau–Yosida regularization: Theoretical preliminaries*, SIAM J. Optim., 7 (1997), pp. 367–385.
- [10] C. LEMARÉCHAL, J. J. STRODIOT AND A. BIHAIN, *On a bundle method for nonsmooth optimization*, in Nonlinear Programming 4, O. L. Mangasarian, R. R. Meyer, and S. M. Robinson, eds., Academic Press, New York, 1981, pp. 245–282.
- [11] O. L. MANGASARIAN AND S. FROMOVITZ, *The Fritz John necessary optimality conditions in the presence of equality and inequality constraints*, J. Math. Anal. Appl., 17 (1967), pp. 37–47.
- [12] A. NEUMAIER, *Interval Methods for Systems of Equations*, Cambridge University Press, Cambridge, 1990.
- [13] B. T. POLYAK, *A general method of solving extremum problems*, Sov. Math., Dokl., 8 (1967), pp. 593–597 (transl. from Dokl. Akad. Nauk SSSR, 174 (1967), pp. 33–36).
- [14] L. QI AND X. CHEN, *A preconditioning proximal Newton method for nondifferentiable convex optimization*, Math. Programming, 76 (1997), pp. 411–429.
- [15] D. F. SHANNO AND E. M. SIMANTIRAKI, *Interior point methods for linear and nonlinear programming*, in The State of the Art in Numerical Analysis, I. S. Duff and G. A. Watson, eds., Clarendon Press, Oxford, 1997, pp. 339–362.