

GLOBAL OPTIMIZATION METHOD FOR SOLVING THE MINIMUM MAXIMAL FLOW PROBLEM*

JUN-YA GOTOH,^a NGUYEN VAN THOAI^{b,*} and YOSHITSUGU YAMAMOTO^{a,†}

^aInstitute of Policy and Planning Sciences, University of Tsukuba, Tsukuba 305-8573, Japan;

^bFB4-Department of Mathematics, University of Trier, 54286 Trier, Germany

(Received 15 October 2002; Revised 5 March 2003; In final form 7 April 2003)

The problem of minimizing the flow value attained by maximal flows plays an important and interesting role to investigate how inefficiently a network can be utilized. It is a typical multiextremal optimization problem, which can have local optima different from global optima. We formulate this problem as a global optimization problem with a special structure and propose a method to combine different techniques in local search and global optimization. Within the proposed algorithm, the advantageous structure of network flow is fully exploited so that the algorithm should be suitable for handling the problem of moderate sizes.

Keywords: Minimum maximal flow; Global optimization; Optimization over efficient sets; Adjacent vertex search; Branch and bound

1 INTRODUCTION

Consider a directed network $N(V, E, s, t, c)$, where V is the set of $m + 2$ nodes, E is the set of n arcs, s is the single source node, t is the single sink node, and c is the vector of arc capacities. A vector x of dimension n is said to be a *feasible flow* if it satisfies the system of conservation equations and capacity constraints:

$$Ax = 0; \quad 0 \leq x \leq c, \quad (1.1)$$

where A is the well-known node–arc incidence matrix restricted to the node set $V \setminus \{s, t\}$, whose size is then $m \times n$. We denote by X the set of feasible flows, i.e.,

$$X = \{x \mid x \in \mathbb{R}^n; Ax = 0; 0 \leq x \leq c\}. \quad (1.2)$$

A vector $x \in X$ is called a *maximal flow* if there does not exist $x' \in X$ such that $x' \geq x$ and $x' \neq x$. We denote the set of all maximal flows by X_M . Further, let $\Delta^+(s)$ and $\Delta^-(s)$ denote

* This paper is written during the second author's stay as a visiting professor at the Institute of Policy and Planning Sciences, University of Tsukuba, Japan.

† Corresponding author. Tel.: +81-29-853-5001/+81-29-853-5202; Fax: +81-29-853-5070; E-mail: yamamoto@sk.tsukuba.ac.jp/dcyama@sk.tsukuba.ac.jp

the sets of arcs leaving and entering the source node s , respectively. Then the total amount of flow, called the *flow value*, of x is given by

$$\sum_{h \in \Delta^+(s)} x_h - \sum_{h \in \Delta^-(s)} x_h. \tag{1.3}$$

The problem to be considered in this article is the minimization of the flow value over the set of all maximal flows, i.e.,

$$P \quad \begin{cases} \min & dx \\ \text{s.t.} & x \in X_M, \end{cases}$$

where d is an n -dimensional row vector defined as

$$d_h = \begin{cases} 1 & \text{if } h \in \Delta^+(s) \\ -1 & \text{if } h \in \Delta^-(s) \\ 0 & \text{otherwise.} \end{cases} \tag{1.4}$$

Problem P was considered in Refs. [27,28] and is closely related to the uncontrollable flow raised by Iri [18,19]. It arises from the following situation. Considering the maximum flow problem, we usually take it for granted that each arc flow is controllable, i.e., we can freely increase and decrease it as long as the conservation equations and capacity constraints are kept satisfied. However, in the situation where we are not able or allowed to reduce the given arc flow, we may fail to reach a *maximum* flow and get stuck in an undesired *maximal* flow. With such restricted controllability, we may end up with different maximal flows depending on the initial flow as well as the way of augmentation. Therefore the minimum of the flow values that are attained by maximal flows will play a prominent role in evaluating how inefficiently the network can be utilized. Note that the problem encompasses the minimum maximal matching problem, which is known to be *NP*-hard, e.g., [14].

Since the set X_M is in general nonconvex, Problem P is one of the typical multiextremal optimization problems. See e.g., Refs. [15,17]. Actually, the set X_M can be considered as the set of all efficient solutions of the multiple objective programming problem (vector optimization problem)

$$MO \quad \begin{cases} \text{vector max} & x \\ \text{s.t.} & x \in X, \end{cases}$$

so that Problem P is a special case of the class of optimization problems over an efficient set. Solution methods for optimization problems over an efficient set can be found e.g., in Refs. [2,4,5,7,8,11,16,20,23,25,30,32,33,36–38]. A common feature of these methods is, however, that they can be only successfully applied to problems where the underlying multiple objective programming problem has a small number of objective functions.

In the present article, we first formulate the underlying problem equivalently as a linear program with an additional nonconvex constraint, and then propose a method to combine local and global optimization techniques for solving the resulting problem in a way that the advantageous network problem structure will be successfully applied.

The equivalent formulation of Problem P is discussed in the next section. Suitable linear program relaxations of this equivalent problem is established in Section 3. Section 4 presents different local and global optimization techniques which are used for the establishment of the combination algorithm in Section 5. The convergence of the combination algorithm depends

on the branching procedure within a branch and bound scheme for checking global optimality. The case of finite convergence is discussed in Section 4 while establishing the branch and bound method using integral rectangular division for the branching procedure. For the use of simplicial branch and bound procedure, some kind of approximate optimal solutions is introduced in Section 5 so that the combination algorithm yields an approximate optimal solution after finitely many iterations. The final section contains an illustrative example and some conclusions.

2 EQUIVALENT PROBLEM FORMULATIONS

In this paper we denote by \mathbb{R}^k and \mathbb{R}_k the set of k -dimensional column vectors and the set of k -dimensional row vectors, respectively. As mentioned in the previous section, the set X_M of maximal flows is exactly the efficient set of MO. From well-known results in multiple objective programming, e.g., Refs. [6,26,29,35], there is a compact subset, say Λ , of $\mathbb{R}_{n++} = \{\lambda \mid \lambda \in \mathbb{R}_n; \lambda > 0\}$ such that a point x belongs to X_M if and only if it maximizes λx over X for some $\lambda \in \Lambda$, i.e.,

$$X_M = \{x \mid x \in X; \lambda x \geq \phi(\lambda) \text{ for some } \lambda \in \Lambda\}, \tag{2.1}$$

where

$$\phi(\lambda) = \max\{\lambda y \mid y \in X\}. \tag{2.2}$$

In what follows we denote by e the row vector of ones and by \mathbb{Z}_n the set of n -dimensional integral row vectors. The following theorem shows that a finite set of integral points of \mathbb{R}_{n++} suffices as Λ .

THEOREM 2.1 $\Lambda_1 = \{\lambda \mid \lambda \in \mathbb{Z}_n; e \leq \lambda \leq ne\}$ suffices as Λ in (2.1).

To prove the above theorem we need the following lemmas.

Let $\bar{x} \in \mathbb{R}^n$ be a given maximal flow. Further let F be the index set defined by $F = \{h \mid h \in E; \bar{x}_h = c_h\}$ and $\bar{F} = E \setminus F$. Note that $F \neq \emptyset$. We refer to a directed path from node i to node j as an i - j path.

LEMMA 2.2 Let G be the graph of node set V and arc set \bar{F} .

- (i) G is acyclic and does not contain an s - t path or a t - s path.
- (ii) For each node $i \in V \setminus \{s, t\}$ at least one of the following two cases occurs:
 - Case 1: G has neither an s - i path nor a t - i path.
 - Case 2: G has neither an i - s path nor an i - t path.

Proof The assertion (i) is clear from the fact that \bar{x} is a maximal flow. Let i be an arbitrary node and suppose that Case 1 of (ii) does not occur, i.e., there is either an s - i path or a t - i path. If there is an s - i path, we have by (i) that there is neither an i - s path nor an i - t path, and if there is a t - i path, we see that there is neither an i - s path nor an i - t path. These correspond to Case 2. ■

Next let a_ℓ denote the row of the incidence matrix A of the network corresponding to node $\ell \in V \setminus \{s, t\}$. Suppose we are given a nonempty subset U of $V \setminus \{s, t\}$ and let

$$\Delta_E^+(U) = \{h \mid h = (i, j) \in E; i \in U; j \in V \setminus U\} \tag{2.3}$$

$$\Delta_E^-(U) = \{h \mid h = (i, j) \in E; j \in U; i \in V \setminus U\}. \tag{2.4}$$

Then it will be readily seen from the definition of the incidence matrix that

$$\sum_{\ell \in U} a_\ell = \sum_{k \in \Delta_E^+(U)} e_k + \sum_{k \in \Delta_E^-(U)} (-e_k), \tag{2.5}$$

where e_k is the k th unit vector of \mathbb{R}_n .

LEMMA 2.3 *For each $h \in \bar{F}$ it holds that*

$$e_h = \alpha_h \sum_{\ell \in V_h} a_\ell + \sum_{k \in F} \beta_{hk} e_k - \sum_{k \in E \setminus \{h\}} \gamma_{hk} e_k \tag{2.6}$$

for some $\alpha_h \in \{-1, 1\}$, $V_h \subseteq V \setminus \{s, t\}$, $\beta_{hk} \in \{0, 1\}$ and $\gamma_{hk} \in \{0, 1\}$.

Proof Let $h = (i, j)$ and consider the following two cases.

Case 1 Node i satisfies the condition of Case 1 of Lemma 2.2.

Let

$$V_h^+ = \{\ell \mid \ell \in V; \text{ there is an } \ell\text{-}i \text{ path of } G\}. \tag{2.7}$$

Then we see from Lemma 2.2 that $s, t, j \notin V_h^+$ and that no arcs of \bar{F} come into V_h^+ from its complement $\overline{V_h^+} = V \setminus V_h^+$. Therefore the cut $(V_h^+, \overline{V_h^+})$ consists of the three sets of arcs: $\Delta_{\bar{F}}^+(V_h^+)$, $\Delta_F^+(V_h^+)$ and $\Delta_{\bar{F}}^-(V_h^+)$. By (2.5) we obtain

$$\sum_{\ell \in V_h^+} a_\ell = \sum_{k \in \Delta_{\bar{F}}^+(V_h^+)} e_k + \sum_{k \in \Delta_F^+(V_h^+)} e_k + \sum_{k \in \Delta_{\bar{F}}^-(V_h^+)} (-e_k), \tag{2.8}$$

which is rewritten as, since $h \in \Delta_{\bar{F}}^+(V_h^+)$,

$$\sum_{\ell \in V_h^+} a_\ell = e_h + \sum_{k \in \Delta_{\bar{F}}^+(V_h^+) \setminus \{h\}} e_k + \sum_{k \in \Delta_F^+(V_h^+)} e_k + \sum_{k \in \Delta_{\bar{F}}^-(V_h^+)} (-e_k). \tag{2.9}$$

Thus we obtain

$$e_h = \sum_{\ell \in V_h^+} a_\ell + \sum_{k \in \Delta_{\bar{F}}^-(V_h^+)} e_k - \left(\sum_{k \in \Delta_F^+(V_h^+)} e_k + \sum_{k \in \Delta_{\bar{F}}^+(V_h^+) \setminus \{h\}} e_k \right). \tag{2.10}$$

Case 2 Node i satisfies the condition of Case 2 of Lemma 2.2.

Since node i satisfies the condition of Case 2 and arc $h = (i, j)$ is in \bar{F} , node j also satisfies that condition. Let $V_h^- = \{\ell \mid \ell \in V; \text{ there is a } j\text{-}\ell \text{ path of } G\}$. Then we see $s, t, i \notin V_h^-$ and that no arcs of \bar{F} go from V_h^- into $\overline{V_h^-} = V \setminus V_h^-$, and the cut $(V_h^-, \overline{V_h^-})$ consists of $\Delta_{\bar{F}}^-(V_h^-)$, $\Delta_F^-(V_h^-)$ and $\Delta_{\bar{F}}^+(V_h^-)$. Therefore

$$\sum_{\ell \in V_h^-} a_\ell = \sum_{k \in \Delta_{\bar{F}}^-(V_h^-)} (-e_k) + \sum_{k \in \Delta_F^-(V_h^-)} (-e_k) + \sum_{k \in \Delta_{\bar{F}}^+(V_h^-)} e_k \tag{2.11}$$

$$= -e_h + \sum_{k \in \Delta_{\bar{F}}^-(V_h^-) \setminus \{h\}} (-e_k) + \sum_{k \in \Delta_F^-(V_h^-)} (-e_k) + \sum_{k \in \Delta_{\bar{F}}^+(V_h^-)} e_k. \tag{2.12}$$

Hence

$$e_h = \left(- \sum_{\ell \in V_h^-} a_\ell \right) + \sum_{k \in \Delta_{\bar{F}}^+(V_h^-)} e_k - \left(\sum_{k \in \Delta_{\bar{F}}^-(V_h^-)} e_k + \sum_{k \in \Delta_{\bar{F}}^-(V_h^-) \setminus \{h\}} e_k \right). \tag{2.13}$$

This completes the proof. ■

Proof of Theorem 2.1 By Lemma 2.3 we see for each $h \in \bar{F}$

$$e_h + \sum_{k \in E \setminus \{h\}} \gamma_{hk} e_k = \alpha_h \sum_{\ell \in V_h} a_\ell + \sum_{k \in F} \beta_{hk} e_k \tag{2.14}$$

for some $\alpha_h \in \{-1, 1\}$, $V_h \subseteq V \setminus \{s, t\}$, $\beta_{hk} \in \{0, 1\}$ and $\gamma_{hk} \in \{0, 1\}$. Adding these equations over $h \in \bar{F}$ and the identities $e_h = e_h$ for $h \in F$, we obtain

$$\sum_{k \in E} \lambda_k e_k = \sum_{\ell \in V \setminus \{s, t\}} \delta_\ell a_\ell + \sum_{k \in F} \zeta_k e_k, \tag{2.15}$$

where $\lambda_k = 1 + \sum_{h \in E \setminus \{k\}} \gamma_{hk}$ for $k \in E$, $\zeta_k = \sum_{h \in F} \beta_{hk} + 1$ for $k \in F$, and δ_ℓ is appropriately defined for $\ell \in V \setminus \{s, t\}$. Note that

$$1 \leq \lambda_k \leq 1 + (n - 1) = n \tag{2.16}$$

for $k \in E$ and $\zeta_k \geq 0$ for $k \in F$. Let $\lambda = \sum_{k \in E} \lambda_k e_k$. Clearly, $\lambda \in \Lambda_1$. Then for any feasible flow x it holds that

$$\lambda \bar{x} = \sum_{k \in E} \lambda_k e_k \bar{x} = \sum_{\ell \in V \setminus \{s, t\}} \delta_\ell a_\ell \bar{x} + \sum_{k \in F} \zeta_k e_k \bar{x} \tag{2.17}$$

$$= \sum_{k \in F} \zeta_k \bar{x}_k = \sum_{k \in F} \zeta_k c_k \tag{2.18}$$

$$\geq \sum_{k \in F} \zeta_k x_k = \sum_{\ell \in V \setminus \{s, t\}} \delta_\ell a_\ell x + \sum_{k \in F} \zeta_k e_k x = \lambda x, \tag{2.19}$$

meaning that the maximal flow \bar{x} maximizes λx over the set of feasible flows. ■

By the property that $\phi(\alpha\lambda) = \alpha\phi(\lambda)$ for $\alpha > 0$, any compact subset of \mathbb{R}_{n++} whose conical hull contains the conical hull of Λ_1 works as Λ . Therefore we could replace Λ by the simplex

$$\Lambda_2 = \left\{ \lambda \mid \lambda \in \mathbb{R}_n; \lambda \geq e; \sum_{i=1}^n \lambda_i = M \right\} \tag{2.20}$$

if M is sufficiently large.

COROLLARY 2.4 n^2 suffices for M defining Λ_2 of (2.20).

Proof Let \bar{x} be a maximal flow. By Theorem 2.1 it maximizes λx over the feasible flows for some $\lambda \in \mathbb{R}_n$ such that $1 \leq \lambda_h \leq n$ for each $h \in E$. Let $\tilde{\lambda} = (n^2 / \sum_{h \in E} \lambda_h) \lambda$. Then since $n^2 \geq \sum_{h \in E} \lambda_h$, $\tilde{\lambda}$ lies in Λ_2 defined for $M = n^2$ and \bar{x} maximizes $\tilde{\lambda} x$ over the feasible flows. ■

Theorem 2.1 implies that Problem P can be solved in theory by the following method. For each integral vector $\lambda \in \Lambda_1$ identify the optimal face $F(\lambda)$ of $\max\{\lambda x \mid x \in X\}$, and solve $\min\{dx \mid x \in F(\lambda)\}$ for a solution $x(\lambda)$. Note that any point of $F(\lambda)$ is a maximal flow, and it is readily seen that $x(\lambda^*)$ such that $dx(\lambda^*) = \min\{dx(\lambda) \mid \lambda \in \Lambda_1\}$ is a solution of P . However, the integral points to be considered amount to n^n , so that this method is not practical.

Remark 2.5 The optimal face $F(\lambda)$ is an *outer semicontinuous mapping* when considered as a point to set mapping, i.e., for a sequence $\{\lambda_\nu\}$ converging to λ any cluster point of the sequence $\{x^\nu\}$ with $x^\nu \in F(\lambda_\nu)$ is contained in $F(\lambda)$. See Exercise 1.19 of Rockafellar and Wets [24]. Then $dx(\lambda)$ is a lower semicontinuous function in λ , i.e., $\liminf_\nu dx(\lambda_\nu) \geq dx(\lambda)$ for any sequence $\{\lambda_\nu\}$ converging to λ . Therefore for a given $\varepsilon > 0$ each λ has a neighborhood such that $dx(\lambda') \geq dx(\lambda) - \varepsilon$ holds for any λ' in the neighborhood. This means that it is very likely that λ 's with large objective function values $dx(\lambda)$ make a cluster. Thus the divide and conquer principle or the branch and bound method should work efficiently.

The above arguments yield two different representations of the set X_M of maximal flows:

$$X_M = \{x \mid x \in X; \lambda x \geq \phi(\lambda) \text{ for some } \lambda \in \Lambda_1\} \tag{2.21}$$

$$= \{x \mid x \in X; \lambda x \geq \phi(\lambda) \text{ for some } \lambda \in \Lambda_2\}, \tag{2.22}$$

each of which will in the following sections provide a scheme for solving the problem. Now Problem P is written equivalently as

$$P \quad \left| \begin{array}{l} \min \quad dx \\ \text{s.t.} \quad x \in X \\ \lambda x - \phi(\lambda) \geq 0 \\ \lambda \in \Lambda, \end{array} \right.$$

where Λ is either Λ_1 or Λ_2 with $M = n^2$.

LEMMA 2.6 *Assume that each arc capacity c_h is a nonnegative integer. Let X_V denote the set of vertices of the polytope X , and let $\bar{x} \in X_V \cap X_M$. Then, whenever $\{x \mid x \in X_M; dx < d\bar{x}\} \neq \emptyset$, there exists $x \in X_M \cap X_V$ such that $dx \leq d\bar{x} - 1$.*

Proof Note first that d is an integral vector. Then this lemma is a direct consequence of the two well-known facts that each vertex of X is an integral vector and that Problem P has an optimal solution in the vertex set of X . ■

3 LINEAR PROGRAM RELAXATION

In this section we explain a linear program relaxation of Problem P . For the sake of further argument, we consider the following problem with λ restricted to a polytope, say S , contained in Λ :

$$P(S) \quad \left| \begin{array}{l} \min \quad dx \\ \text{s.t.} \quad Ax = 0 \\ 0 \leq x \leq c \\ \lambda x - \phi(\lambda) \geq 0 \\ \lambda \in S. \end{array} \right.$$

Let

$$\Phi(S) = \min\{\phi(\lambda) \mid \lambda \in S\}. \tag{3.1}$$

Our method for constructing linear program relaxations of Problem $P(S)$ is based on the following result.

LEMMA 3.1 *Let $\{\lambda_1, \dots, \lambda_q\}$ be the vertex set of S . Then Problem $P(S)$ is relaxed to the following linear program in variables $x^\ell \in \mathbb{R}^n$ for $\ell = 1, \dots, q$:*

$$\bar{P}(S) \quad \left| \begin{array}{l} \min \quad \sum_{\ell=1}^q dx^\ell \\ \text{s.t.} \quad Ax^\ell = 0 \quad \text{for } \ell = 1, \dots, q \\ \quad \quad x^\ell \geq 0 \quad \text{for } \ell = 1, \dots, q \\ \quad \quad \sum_{\ell=1}^q x^\ell \leq c \\ \quad \quad \sum_{\ell=1}^q \lambda_\ell x^\ell - \Phi(S) \geq 0, \end{array} \right.$$

i.e., the optimal value $\bar{\mu}(S)$ of Problem $\bar{P}(S)$ yields a lower bound of the optimal value of Problem $P(S)$.

Proof We show that for any feasible solution (x, λ) of Problem $P(S)$ there exists a feasible solution (x^1, \dots, x^q) of Problem $\bar{P}(S)$ satisfying

$$dx = \sum_{\ell=1}^q dx^\ell. \tag{3.2}$$

Since $\lambda \in S$,

$$\lambda = \sum_{\ell=1}^q \beta_\ell \lambda_\ell \tag{3.3}$$

for some nonnegative numbers β_ℓ ($\ell = 1, \dots, q$) such that $\sum_{\ell=1}^q \beta_\ell = 1$. Define

$$x^\ell = \beta_\ell x \quad \text{for } \ell = 1, \dots, q. \tag{3.4}$$

Then clearly $\sum_{\ell=1}^q x^\ell = x$ and (x^1, \dots, x^q) satisfies the first three constraints of $\bar{P}(S)$. For the last constraint we have

$$\sum_{\ell=1}^q \lambda_\ell x^\ell = \sum_{\ell=1}^q \lambda_\ell \beta_\ell x = \lambda x \geq \phi(\lambda) \geq \Phi(S), \tag{3.5}$$

where the last two inequalities follow from the assumption that (x, λ) is feasible to $P(S)$ and from the definition of $\Phi(S)$ in (3.1). ■

Remark 3.2 If Problem $\bar{P}(S)$ is infeasible, so is Problem $P(S)$. In this case we set $\bar{\mu}(S) = +\infty$.

Remark 3.3 The value $\Phi(S)$ is determined in two different ways, both of which come from the bilinearity of λx . The first way is

$$\begin{aligned}\Phi(S) &= \min_{\lambda \in S} \max_{x \in X} \lambda x \\ &= \min\{r \mid \lambda \in S; r \geq \lambda x \text{ for all } x \in X\} \\ &= \min\{r \mid \lambda \in S; r \geq \lambda x^\ell \text{ for all } x^\ell \in X_V\}.\end{aligned}$$

Note that we need not know all vertices of X in advance if we use the so-called column generation technique in solving this problem. Applying the well-known minimax theorem yields the second way:

$$\begin{aligned}\Phi(S) &= \min_{\lambda \in S} \max_{x \in X} \lambda x \\ &= \max_{x \in X} \min_{\lambda \in S} \lambda x \\ &= \max\{r \mid x \in X; r \leq \lambda x \text{ for all } \lambda \in S\} \\ &= \max\{r \mid x \in X; r \leq \lambda_\ell x \text{ for } \ell = 1, \dots, q\}.\end{aligned}$$

In either way $\Phi(S)$ is obtained by solving a linear programming.

Remark 3.4 The relaxation problem $\bar{P}(S)$ is a multicommodity flow problem with a side constraint, to which various techniques for the multicommodity flow problem can be applied, e.g., Lagrangian relaxation, column generation approach or Dantzig-Wolfe decomposition, resource-directive decomposition, and basis partitioning method in Ref. [1]. Here we explain first the column generation approach, and then the Lagrangian relaxation technique.

Let $Y = \{x \mid x \in \mathbb{R}_+^n; Ax = 0\}$, then Y is a pointed convex cone, and is a conical hull of a finite number of nonzero vectors, say $g^1, \dots, g^r \in \mathbb{R}_+^n$. Let G be the $n \times r$ matrix of these column vectors, then each $x \in Y$ is written as $x = G\alpha$ for some $\alpha \in \mathbb{R}_+^r$. Problem $\bar{P}(S)$ is then

$$\left. \begin{array}{l} \min \quad \sum_{\ell=1}^q d G \alpha^\ell \\ \text{s.t.} \quad \sum_{\ell=1}^q G \alpha^\ell \leq c \\ \quad \quad \sum_{\ell=1}^q \lambda_\ell G \alpha^\ell - \Phi(S) \geq 0 \\ \quad \quad \alpha^\ell \geq 0 \quad \text{for } \ell = 1, \dots, q. \end{array} \right\}$$

Suppose we have a simplex multiplier vector $(\pi, \pi_0) \in \mathbb{R}_{n+1}$ corresponding to a feasible basis, and search for a nonbasic variable to be pivoted in. Noting the reduced cost of the nonbasic variable α_i^ℓ is given by

$$d g^i - (-\pi g^i + \pi_0(\lambda_\ell g^i)) = (d + \pi - \pi_0 \lambda_\ell) g^i,$$

the problem to be solved is

$$\min_{\ell=1, \dots, q} \min_{i=1, \dots, r} (d + \pi - \pi_0 \lambda_\ell) g^i = \min_{\ell=1, \dots, q} \min_{y \in Y} (d + \pi - \pi_0 \lambda_\ell) y.$$

Note that for each ℓ , $\min_{y \in Y} (d + \pi - \pi_0 \lambda_\ell)y$ is a shortest path and cycle problem. When the minimum value above is nonnegative, the current basic solution is optimal to $\bar{P}(S)$.

Next consider the Lagrangian relaxation problem with a multiplier $\tau \geq 0$

$$\left| \begin{array}{l} \min \quad \sum_{\ell=1}^q dx^\ell + \tau \left(\Phi(S) - \sum_{\ell=1}^q \lambda_\ell x^\ell \right) \\ \text{s.t.} \quad Ax^\ell = 0 \quad \text{for } \ell = 1, \dots, q \\ \quad \quad x^\ell \geq 0 \quad \text{for } \ell = 1, \dots, q \\ \quad \quad \sum_{\ell=1}^q x^\ell \leq c. \end{array} \right.$$

For each $j = 1, 2, \dots, n$ let

$$\lambda_j^{\max} = \max\{(\lambda_\ell)_j \mid \ell = 1, \dots, q\}$$

and let $d'(\tau) \in \mathbb{R}_n$ be the vector whose j th component $d'_j(\tau)$ is defined by

$$d'_j(\tau) = d_j - \tau \lambda_j^{\max}.$$

Then clearly for a feasible solution (x^1, \dots, x^q)

$$\sum_{\ell=1}^q d'(\tau)x^\ell \leq \sum_{\ell=1}^q (d - \tau \lambda_\ell)x^\ell,$$

so that the above Lagrangian relaxation can be relaxed further by the problem

$$\left| \begin{array}{l} \min \quad \sum_{\ell=1}^q d'(\tau)x^\ell + \tau \Phi(S) \\ \text{s.t.} \quad Ax^\ell = 0 \quad \text{for } \ell = 1, \dots, q \\ \quad \quad x^\ell \geq 0 \quad \text{for } \ell = 1, \dots, q \\ \quad \quad \sum_{\ell=1}^q x^\ell \leq c, \end{array} \right.$$

which is actually equivalent to the following minimum cost network flow problem in n variables:

$$\bar{\bar{P}}(S) \quad \left| \begin{array}{l} \min \quad d'(\tau)x + \tau \Phi(S) \\ \text{s.t.} \quad Ax = 0 \\ \quad \quad 0 \leq x \leq c. \end{array} \right.$$

An efficient choice for the multiplier τ can be taken using the well-known parametric linear programming technique. See for example Refs. [1,3,10].

The dual problem of $\bar{P}(S)$, denoted by $\bar{D}(S)$, is written as

$$\bar{D}(S) \quad \left| \begin{array}{l} \max \quad -vc + z\Phi(S) \\ \text{s.t.} \quad u_\ell A - v + z\lambda_\ell \leq d \quad \text{for } \ell = 1, \dots, q \\ \quad \quad (v, z) \geq 0, \end{array} \right.$$

where $u_k \in \mathbb{R}_m, v \in \mathbb{R}_n, z \in \mathbb{R}$.

LEMMA 3.5 A nonnegative vector (v, z) satisfies the constraints of $\bar{D}(S)$ together with some $u_1, \dots, u_q \in \mathbb{R}_m$ if and only if for any $\lambda \in S$ there is $u(\lambda) \in \mathbb{R}_m$ such that $u(\lambda)A - v + z\lambda \leq d$.

Proof The “if” part is readily seen by setting $u_\ell = u(\lambda_\ell)$ for $\ell = 1, \dots, q$. To show the “only if” part, let λ be a given vector of S . Then $\lambda = \sum_{\ell=1}^q \beta_\ell \lambda_\ell$ for some nonnegative β_ℓ 's with $\sum_{\ell=1}^q \beta_\ell = 1$. Let $u(\lambda) = \sum_{\ell=1}^q \beta_\ell u_\ell$. Then

$$\begin{aligned} u(\lambda)A - v + z\lambda &= \left(\sum_{\ell=1}^q \beta_\ell u_\ell \right) A - v + z \left(\sum_{\ell=1}^q \beta_\ell \lambda_\ell \right) \\ &= \sum_{\ell=1}^q \beta_\ell (u_\ell A - v + z\lambda_\ell) \leq d. \end{aligned} \quad \blacksquare$$

The lower bound $\bar{\mu}(S)$ has the following properties, which will be utilized within the branch and bound procedure.

LEMMA 3.6 Suppose $\emptyset \neq S_2 \subseteq S_1 \subseteq \Lambda$. Then

$$-\infty < \bar{\mu}(S_1) \leq \bar{\mu}(S_2). \tag{3.6}$$

Proof Suppose (v, z) is a feasible solution of $\bar{D}(S_1)$ together with (u_1, \dots, u_q) . Then by Lemma 3.5, we see that (v, w, z) is a feasible solution of $\bar{D}(S_2)$ with some (u'_1, \dots, u'_q) . Since the objective function value of $\bar{D}(S)$ is determined solely by (v, z) , we see the monotonicity $\bar{\mu}(S_1) \leq \bar{\mu}(S_2)$. The fact that $-\infty < \bar{\mu}(S_1)$ is a direct consequence of the boundedness of the feasible region of $\bar{P}(S_1)$. ■

We now show that the relaxation problem $\bar{P}(S)$ can be substantially simplified when S is a hyper rectangle.

LEMMA 3.7 If $\lambda \leq \lambda'$, then $\phi(\lambda) \leq \phi(\lambda')$.

Proof Let \bar{x} be a point of X that maximizes λx over X . Since $\bar{x} \geq 0$ we have $(\lambda - \lambda')\bar{x} \leq 0$, and hence

$$\phi(\lambda) = \lambda\bar{x} \leq \lambda'\bar{x} \leq \phi(\lambda'). \tag{3.7}$$

LEMMA 3.8 Let $\{\lambda_1, \dots, \lambda_q\}$ be the vertex set of $S \subseteq \Lambda$ and suppose $\lambda_1 \geq \lambda_\ell \geq \lambda_q$ for all $\ell = 1, \dots, q$. Then Problem $\bar{P}(S)$ is equivalent to the problem

$$\bar{P}'(S) \quad \left\{ \begin{array}{l} \min \quad dx \\ \text{s.t.} \quad Ax = 0 \\ \quad \quad 0 \leq x \leq c \\ \quad \quad \lambda_1 x - \phi(\lambda_q) \geq 0 \end{array} \right.$$

in n variables.

Proof Note that $\lambda_q \leq \lambda_\ell$ for all $\ell = 1, \dots, q$ implies that $\lambda_q \leq \lambda$ for all $\lambda \in S$. Then by Lemma 3.7 we have $\Phi(S) = \phi(\lambda_q)$.

Let (x^1, \dots, x^q) be a feasible solution of $\bar{P}(S)$ and let $x = \sum_{\ell=1}^q x^\ell$. Then clearly $Ax = 0$ and $0 \leq x \leq c$. Furthermore, since $x^\ell \geq 0$ we obtain

$$\begin{aligned} \lambda_1 x &= \lambda_1 \sum_{\ell=1}^q x^\ell = \sum_{\ell=1}^q \lambda_1 x^\ell \geq \sum_{\ell=1}^q \lambda_\ell x^\ell \geq \Phi(S) \\ dx &= \sum_{\ell=1}^q dx^\ell. \end{aligned}$$

This means that $\bar{\mu}(S) \geq \bar{\mu}'(S)$, where $\bar{\mu}'(S)$ is the optimal objective function value of $\bar{P}'(S)$. Let x be a feasible solution of $\bar{P}'(S)$ and let $x^1 = x, x^2 = \dots = x^q = 0$. Then (x^1, \dots, x^q) clearly satisfies the constraints of $\bar{P}(S)$, meaning $\bar{\mu}(S) \leq \bar{\mu}'(S)$. ■

From Lemma 3.8 if we take Λ_1 as Λ and consider a hyper rectangle $S = \{\lambda \mid \lambda \in \Lambda_1; \underline{\lambda} \leq \lambda \leq \bar{\lambda}\}$ contained in Λ_1 , then, although S has as many as 2^n vertices, the relaxation problem $\bar{P}'(S)$ has only n variables as Problem $P(S)$ does. This might be an advantage of using Λ_1 .

However, the simplicity of $\bar{P}'(S)$ can be to its disadvantage because $\lambda x - \phi(\underline{\lambda}) \geq 0$ is a too much relaxed constraint of $\lambda x - \phi(\lambda) \geq 0$. Therefore we propose below an improvement with aid of the concave envelope of the bilinear term λx and an approximation of the function $\phi(\lambda)$.

LEMMA 3.9 Let $S = \{\lambda \mid \lambda \in \Lambda_1; \underline{\lambda} \leq \lambda \leq \bar{\lambda}\}$, and let V be a nonempty subset of $X_M \cap X_V$. Then the problem

$$\bar{P}''(S, V) \quad \left| \begin{array}{l} \min \quad dx \\ \text{s.t.} \quad Ax = 0 \\ \quad \quad 0 \leq x \leq c \\ \quad \quad g_k^1(\lambda_k, x_k) - t_k \geq 0 \quad \text{for } k \in E \\ \quad \quad g_k^2(\lambda_k, x_k) - t_k \geq 0 \quad \text{for } k \in E \\ \quad \quad \sum_{k \in E} t_k - \lambda v \geq 0 \quad \text{for } v \in V \end{array} \right.$$

is a relaxation problem of $P(S)$, where

$$\begin{aligned} g_k^1(\lambda_k, x_k) &= c_k \lambda_k + \underline{\lambda}_k x_k - \underline{\lambda}_k c_k \\ g_k^2(\lambda_k, x_k) &= \bar{\lambda}_k x_k. \end{aligned}$$

Proof First note that $\bar{P}''(S, V)$ is equivalent to

$$\left| \begin{array}{l} \min \quad dx \\ \text{s.t.} \quad Ax = 0 \\ \quad \quad 0 \leq x \leq c \\ \quad \quad \sum_{k \in E} \min\{g_k^1(\lambda_k, x_k), g_k^2(\lambda_k, x_k)\} \geq \max\{\lambda v \mid v \in V\}. \end{array} \right.$$

The left hand side $\sum_{k \in E} \min\{g_k^1(\lambda_k, x_k), g_k^2(\lambda_k, x_k)\}$ of the last constraint is the concave envelope of λx on the set $S \times \{x \mid x \in \mathbb{R}^n; 0 \leq x \leq c\}$ (see, for example, Corollary 3.6 of [34] and Theorem IV.7 of [17]), and hence

$$\sum_{k \in E} \min\{g_k^1(\lambda_k, x_k), g_k^2(\lambda_k, x_k)\} \geq \lambda x \tag{3.8}$$

holds on this set. On the other hand, since $V \subseteq X_M \cap X_V$,

$$\phi(\lambda) = \max\{\lambda x \mid x \in X\} \geq \max\{\lambda v \mid v \in V\}. \tag{3.9}$$

Then combining (3.8) and (3.9) yields the assertion. ■

Remark 3.10 When V contains a vertex, say \underline{v} such that $\underline{\lambda} \underline{v} = \phi(\underline{\lambda})$, the constraints of $\bar{P}''(S, V)$ implies the constraint $\bar{\lambda} x - \phi(\underline{\lambda}) \geq 0$ in the relaxation $\bar{P}'(S)$. In fact, let $(\lambda, x, t) \in \mathbb{R}_n \times \mathbb{R}^n \times \mathbb{R}^n$ be feasible to $\bar{P}''(S, V)$. Then $\sum_{k \in E} t_k \geq \lambda \underline{v}$, implying $\sum_{k \in E} t_k \geq \underline{\lambda} \underline{v} = \phi(\underline{\lambda})$. On the other hand $\bar{\lambda} x = \sum_{k \in E} g_k^2(\lambda_k, x_k) \geq \sum_{k \in E} t_k$. Then we see $\bar{\lambda} x - \phi(\underline{\lambda}) \geq 0$ is satisfied.

The algorithm to be proposed contains a local search procedure, Section 4.1, in which we generate a series of extreme maximal flows, that is, points of $X_M \cap X_V$. By keeping these, we can make the set V at no extra cost. However, it is recommended to add \underline{v} and as well as the point \bar{v} such that $\bar{\lambda} \bar{v} = \phi(\bar{\lambda})$ to the set V .

Remark 3.11 For an incumbent value α let

$$\begin{aligned} \ell_k &= \min\{x_k \mid Ax = 0; 0 \leq x \leq c; dx \leq \alpha\} \\ u_k &= \max\{x_k \mid Ax = 0; 0 \leq x \leq c; dx \leq \alpha\} \end{aligned}$$

for each $k \in E$. Then we have only to consider feasible flows satisfying $\ell_k \leq x_k \leq u_k$ for all $k \in E$. Hence we could replace $0 \leq x_k \leq c_k$ in the above argument by $\ell_k \leq x_k \leq u_k$ to obtain

$$\begin{aligned} g_k^1(\lambda_k, x_k) &= u_k \lambda_k + \underline{\lambda}_k x_k - \underline{\lambda}_k u_k \\ g_k^2(\lambda_k, x_k) &= \ell_k \lambda_k + \bar{\lambda}_k x_k - \bar{\lambda}_k \ell_k, \end{aligned}$$

which might strengthen the relaxation.

4 LOCAL SEARCH AND CHECKING UP GLOBAL OPTIMALITY

4.1 Local Search

The algorithms for the optimization over the efficient set proposed by Philip [22], Ecker and Song [12], Fülöp [13] and Bolintineanu [9] are mainly based on the technique of moving from an efficient vertex to an efficient neighbor with a smaller objective function value via an efficient edge. In this section, following their argument we will explain a local search technique called *Adjacent Vertex Search Procedure*.

We say that a maximal flow is *extreme* if it is a vertex of the set of feasible flows X . It is known, e.g. in Refs. [21,26,35], the set of maximal flows X_M is connected, and all extreme

maximal flows are connected by paths of edges consisting of maximal flows. Thus, starting from an extreme maximal flow, we could reach an optimal solution of Problem P by a series of pivot operations in theory. However, we cannot decrease the objective function value monotonically along the path as we trace, i.e., we might be eventually caught by a non-optimal extreme maximal flow none of whose neighboring extreme maximal flows have a smaller objective function value. When this occurs, it is a local minimum solution. See for example Ref. [9].

The Adjacent Vertex Search (AVS) Procedure goes as follows. We denote by $[x, x']$ the edge of X connecting x and x' for $x, x' \in X_V$ and

$$N_M(x) = \{x' \mid x' \in X_V \cap X_M; [x, x'] \subseteq X_M\}.$$

Adjacent Vertex Search (AVS) Procedure

$\langle\langle$ Initialization $\rangle\rangle$

Find $x^0 \in X_V \cap X_M$. If $N_M(x^0) = \emptyset$, then x^0 is an optimal solution of P . Otherwise, set $k \leftarrow 0$ and go to Step k .

$\langle\langle$ Step k $\rangle\rangle$

$\langle k1$ \rangle If $\{x \mid x \in N_M(x^k); dx < dx^k\} \neq \emptyset$, choose x^{k+1} from this set, let $k \leftarrow k + 1$ and go to Step k .

$\langle k2$ \rangle Otherwise, set $v \leftarrow x^k$ and stop.

Note that the initial extreme maximal flow x^0 is easily found by choosing an arbitrary positive vector λ and maximizing λx over X . The AVS Procedure generates a sequence of distinct extreme maximal flows x^0, x^1, \dots, x^k with decreasing objective function values, which implies owing to the integrality property that $dx^k \leq dx^0 - k$. When the relaxation problem $\bar{P}''(S, V)$ is used in the branch and bound procedure, we accumulate these extreme maximal flows as the initial set V .

4.2 Checking Up Global Optimality

We present in this subsection two *Branch and Bound (BB)* Procedures for handling the following problem:

$$\text{CGO}(\alpha) \quad \left\{ \begin{array}{l} \text{For a given integer } \alpha, \text{ find an extreme maximal flow} \\ v \in X_M \cap X_V \text{ such that } dv \leq \alpha, \\ \text{or show that there does not exist such a point.} \end{array} \right.$$

As shown in Section 2, for the description of the set X_M , we can use one of two sets Λ_1 and Λ_2 . The set Λ_1 defined in Theorem 2.1 consists of all integral vectors contained in the rectangle $\{\lambda \mid \lambda \in \mathbb{R}_n; e \leq \lambda \leq ne\}$, while the set Λ_2 is an $(n - 1)$ -simplex defined in (2.20). Before presenting two branch and bound procedures for handling Problem $\text{CGO}(\alpha)$ according to Λ_1 and Λ_2 , respectively, we propose here two kinds of division process called *integral rectangular division* and *simplicial division*.

4.2.1 Integral Rectangular Division

Let $S \subseteq \Lambda_1$ be a rectangle with integral bound vectors, which contains more than one integral vector and is defined by

$$S = \{\lambda \mid \lambda \in \mathbb{R}_n; \underline{\lambda} \leq \lambda \leq \bar{\lambda}\},$$

where $\underline{\lambda}$ and $\bar{\lambda}$ are integral vectors of \mathbb{R}_n such that $\underline{\lambda} \leq \bar{\lambda}$. Further, let S_1, \dots, S_q be rectangles with integral bound vectors having the following properties:

$$\begin{aligned} \bigcup_{i=1}^q S_i &\subseteq S, \\ S_i \cap S_j &= \emptyset \quad \text{for } i \neq j, \\ \bigcup_{i=1}^q (S_i \cap \mathbb{Z}_n) &= S \cap \mathbb{Z}_n. \end{aligned}$$

Then we say that $\{S_1, \dots, S_q\}$ is an *integral rectangular division* of the rectangle S .

As a special case of this division, we consider the following *integral rectangular bisection*. Let $\lambda' \in S$ such that $\lambda' \neq \bar{\lambda}$, and let $\ell \in \{1, \dots, n\}$ such that $\lambda'_\ell < \bar{\lambda}_\ell$, (note that such an index ℓ exists, whenever S contains more than one integral vector). For each real number t , denoting by $\lfloor t \rfloor$ the largest integer which is less than or equal to t , we define two rectangles

$$\begin{aligned} S_1 &= \{\lambda \mid \lambda \in S; \lambda_\ell \leq \lfloor \lambda'_\ell \rfloor\} \\ S_2 &= \{\lambda \mid \lambda \in S; \lambda_\ell \geq \lfloor \lambda'_\ell \rfloor + 1\}. \end{aligned}$$

Then, it is easy to verify that $\{S_1, S_2\}$ is an integral rectangular division of S . We say that S is divided into $\{S_1, S_2\}$ by an integral rectangular bisection using the point λ' .

4.2.2 *Simplicial Division*

Let S be an $(n - 1)$ -simplex with vertex set $S_V = \{\lambda_1, \dots, \lambda_n\}$. Choose a point $\lambda' \in S \setminus S_V$, which is uniquely represented as

$$\lambda' = \sum_{i=1}^n \beta_i \lambda_i, \quad \beta_i \geq 0 \quad (i = 1, \dots, n), \quad \sum_{i=1}^n \beta_i = 1.$$

For each i such that $\beta_i > 0$, construct the simplex S_i obtained from S by replacing the vertex λ_i by λ' , i.e., $S_i = \text{co}\{\lambda_1, \dots, \lambda_{i-1}, \lambda', \lambda_{i+1}, \dots, \lambda_n\}$, where $\text{co}A$ denotes the convex hull of a set A . This division is called a *radial simplicial division*.

When λ' is the midpoint of a longest edge of S , then we obtain two subsimplices. This special case is called a *simplicial bisection*.

As discussed in the preceding section, our BB Procedures are based on the linear relaxation $\bar{P}(S)$ or $\bar{P}''(S)$ of the subproblem $P(S)$ with λ restricted to a subset $S \subseteq \Lambda$, where Λ is either Λ_1 or Λ_2 . The branching process subdivides Λ into finitely many subsets yielding a class of subproblems to be solved. In the algorithm to be proposed we repeatedly apply the AVS Procedure, which provides a local minimum incumbent solution v^v , and then one of the BB Procedures to check up the global optimality of v^v . The chosen BB procedure starts with the number $\alpha = dv^v - 1$ and the class \mathcal{R} of subsets S of Λ such that $\bar{\mu}(S) \leq \alpha$, and also with the set V of extreme maximal flows obtained so far. In the BB1 we denote the optimal value of the relaxation problem $\bar{P}''(S, V)$ by $\bar{\mu}(S)$ for the simplicity of notation.

Branch and Bound Procedure (BB1) (according to Λ_1)

$\langle\langle$ Initialization $\rangle\rangle$

Set $k \leftarrow 0$ and $\mathcal{R}_0 \leftarrow \mathcal{R}$.

$\langle\langle$ Step k $\rangle\rangle$

- (k1) Set $\mu_k \leftarrow \min\{\bar{\mu}(S) \mid S \in \mathcal{R}_k\}$ and choose $S_k \in \mathcal{R}_k$ such that $\bar{\mu}(S_k) = \mu_k$. Divide S_k into S_{k1}, \dots, S_{kp} by an integral rectangular division and set $\mathcal{R}_k \leftarrow \mathcal{R}_k \setminus \{S_k\} \cup \{S_{k1}, \dots, S_{kp}\}$,
- (k2) For $j = 1, \dots, p$ do:
- (a) If $|S_{kj} \cap \mathbb{Z}_n| = 1$, then do:
 - (i) Set $\mathcal{R}_k \leftarrow \mathcal{R}_k \setminus \{S_{kj}\}$.
 - (ii) Choose $\lambda \in S_{kj} \cap \mathbb{Z}_n$ and solve $\max\{\lambda x \mid x \in X\}$, yielding the optimal face $F(\lambda)$.
 - (iii) Solve $\min\{dx \mid x \in F(\lambda)\}$, yielding a vertex solution $x(\lambda)$.
 - (iv) Set $V \leftarrow V \cup \{x(\lambda)\}$.
 - (v) If $dx(\lambda) \leq \alpha$, then set $w \leftarrow x(\lambda)$ and go to (k4). Otherwise, go to Endfor.
 - (b) If $|S_{kj} \cap \mathbb{Z}_n| > 1$, then do:
 - (i) Solve $P''(S_{kj}, V)$, yielding the optimal value $\bar{\mu}(S_{kj})$ and an optimal solution y if feasible ($\bar{\mu}(S_{kj}) = +\infty$ when infeasible).
 - (ii) If $\bar{\mu}(S_{kj}) > \alpha$, then set $\mathcal{R}_k \leftarrow \mathcal{R}_k \setminus \{S_{kj}\}$ and go to Endfor.
 - (iii) If $y \in X_M$, then identify the minimal face F of X containing y , solve $\min\{dx \mid x \in F\}$ for a vertex solution w and set $V \leftarrow V \cup \{w\}$.
 - (iv) Go to (k4).
- Endfor
- (k3) If $\mathcal{R}_k \neq \emptyset$, then set $\mathcal{R}_{k+1} \leftarrow \mathcal{R}_k, k \leftarrow k + 1$ and go to (Step k). Otherwise, set $\mathcal{R} \leftarrow \emptyset$ and quit.
- (k4) Set $\mathcal{R} \leftarrow \{S \mid S \in \mathcal{R}_k; \bar{\mu}(S) \leq dw - 1\}$ and quit.

THEOREM 4.1 *Procedure BB1 terminates after finitely many iterations, either yielding an extreme maximal flow with an objective function value being less than or equal to α , or indicating that such a maximal flow does not exist. In other words, Procedure BB1 solves Problem CGO(α) finitely.*

Proof Each subsequence of rectangles, $\{S_q\}$, generated throughout Procedure (BB1) such that $S_{q+1} \subset S_q$ for all q , must be finite, since every S_q contains at least one element from \mathbb{Z}_n .

If the procedure terminates at Step (k4), then the point w is an extreme maximal flow satisfying $dw \leq \alpha$.

If the procedure terminates at Step (k3), i.e., $\mathcal{R} = \emptyset$, then it follows that each subset S of Λ_1 yields a lower bound $\bar{\mu}(S) > \alpha$, which implies that there does not exist an extreme maximal flow $v \in X_M \cap X_V$, such that $dv \leq \alpha$. ■

Branch and Bound Procedure (BB2) (according to Λ_2)

((Initialization))

Set $k \leftarrow 0$ and $\mathcal{R}_0 \leftarrow \mathcal{R}$.

((Step k))

- (k1) (a) Set $\mu_k \leftarrow \min\{\bar{\mu}(S) \mid S \in \mathcal{R}_k\}$ and choose $S_k \in \mathcal{R}_k$ such that $\bar{\mu}(S_k) = \mu_k$.
- (b) Divide S_k into S_{k1}, \dots, S_{kp} by a simplicial division and set $\mathcal{R}_k \leftarrow \mathcal{R}_k \setminus \{S_k\} \cup \{S_{k1}, \dots, S_{kp}\}$,
- (k2) For $j = 1, \dots, p$ do:
- (a) Solve $\bar{P}(S_{kj})$, yielding the optimal value $\bar{\mu}(S_{kj})$ and an optimal solution $(x_{kj}^1, \dots, x_{kj}^n)$ if feasible ($\bar{\mu}(S_{kj}) = +\infty$ when infeasible).
 - (b) If $\bar{\mu}(S_{kj}) > \alpha$, then set $\mathcal{R}_k \leftarrow \mathcal{R}_k \setminus \{S_{kj}\}$.
 - (c) Let $x_{kj} \leftarrow \sum_{\ell=1}^n x_{kj}^\ell$.

- (d) If $x_{kj} \in X_M$, then set $y \leftarrow x_{kj}$ and go to $\langle k4 \rangle$.
Endfor
- $\langle k3 \rangle$ (a) If $\mathcal{R}_k \neq \emptyset$, then set $\mathcal{R}_{k+1} \leftarrow \mathcal{R}_k, k \leftarrow k + 1$ and go to $\langle k1 \rangle$.
(b) Otherwise, set $\mathcal{R} \leftarrow \emptyset$ and quit.
- $\langle k4 \rangle$ (a) Identify the minimal face F of X containing y and solve

$$\min\{dx \mid x \in F\}$$

for a vertex solution w .

- (b) Set

$$\mathcal{R} \leftarrow \{S \mid S \in \mathcal{R}_k; \bar{\mu}(S) \leq dw - 1\},$$

and quit.

If the Procedure BB2 terminates after finitely many iterations, then it either yields an extreme maximal flow with an objective function value being less than or equal to α , or indicates that such a maximal flow does not exist when $\mathcal{R} = \emptyset$.

In the case that the procedure is infinite, it generates at least an infinite nested subsequence $\{S_\kappa\}$ of subsimplices such that $S_{\kappa+1} \subset S_\kappa$ for all κ . Below we show that the procedure will generate an infinite sequence of flows converging to a minimum maximal flow, i.e., an optimal solution of Problem P .

We recall the *exhaustiveness* of the division process introduced for the establishment of convergence properties of branch and bound algorithms in global optimization (see, e.g., [31]). An infinite nested sequence of subsets $\{S_\kappa\}_\kappa$ is said to be *exhaustive* if $\bigcap_{\kappa=1}^\infty S_\kappa$ is a singleton. A simplicial division process is called *exhaustive* if each nested infinite subsequence of subsets generated by it is exhaustive. It is well known that the simplicial bisection process is exhaustive.

THEOREM 4.2 *Assume that the division process is exhaustive and Procedure BB2 is infinite. For each k let $x_k = \sum_{\ell=1}^n x_k^\ell$, where (x_k^1, \dots, x_k^n) is an optimal solution to $\bar{P}(S_k)$. Then the sequence $\{x_k\}$ has an accumulation point, and each of them is an optimal solution of Problem P .*

Proof Since $x_k \in X$ for each k and X is a compact set, the sequence $\{x_k\}$ has an accumulation point in X . Let x^* be an arbitrary accumulation point of $\{x_k\}$, and let $\{x_\kappa\}$ be a subsequence converging to x^* . From Lemma 3.5 and the property that $\{\mu_\kappa\}$ is nondecreasing and bounded from above by α , it follows that there exists a limit μ^* of $\{\mu_\kappa\}$. By using subsequences if necessary, assume that $\mu_\kappa \rightarrow \mu^*$ as $\kappa \rightarrow \infty$, and $\{S_\kappa\}$ is the corresponding subsequence of simplices such that $S_{\kappa+1} \subset S_\kappa$ for all κ . Since the simplicial division is exhaustive, it follows that

$$\bigcap_{\kappa=1}^\infty S_\kappa = \{\lambda^*\}, \tag{4.1}$$

and hence, denoting by λ_κ^ℓ ($\ell = 1, \dots, n$) the vertices of S_κ we have $\lambda_\kappa^\ell \rightarrow \lambda^* \in \Lambda_2$ as $\kappa \rightarrow \infty$ for $i = 1, \dots, n$. Thus, we have $x^* \in X, \lambda^* \in \Lambda_2$, and $\lambda^* x^* - \phi(\lambda^*) \geq 0$, which implies that (x^*, λ^*) is a feasible solution of Problem $P(\Lambda_2)$, i.e. Problem P , and therefore, an optimal solution of this problem with the optimal value $dx^* = \mu^*$. ■

5 GLOBAL OPTIMIZATION ALGORITHM AND APPROXIMATE OPTIMAL SOLUTION

Combining Procedure AVS with Procedure BB1 or BB2 in Section 4, we propose the following algorithm for globally solving Problem P .

Global Optimization Algorithm (GOA)

$\langle\langle$ Initialization $\rangle\rangle$

Compute an extreme maximal flow $w^0 \in X_V \cap X_M$. If $N_M(w^0) = \emptyset$, then w^0 is an optimal solution of P . Otherwise, set $\nu \leftarrow 1$, $\mathcal{R} \leftarrow \{\Lambda\}$ and go to Iteration ν .

$\langle\langle$ Iteration ν $\rangle\rangle$

$\langle\nu 1\rangle$ Apply AVS to Problem P starting from $w^{\nu-1}$ and let v^ν be the extreme maximal flow obtained. Set $\alpha_\nu \leftarrow dv^\nu - 1$ and go to Step $\nu 2$.

$\langle\nu 2\rangle$ Set $\mathcal{R} \leftarrow \{S \mid S \in \mathcal{R}; \bar{\mu}(S) \leq \alpha_\nu\}$ and apply BB.

(a) If an extreme maximal flow w^ν with $dw^\nu \leq \alpha_\nu$ is found, set $\nu \leftarrow \nu + 1$ and go to Iteration ν .

(b) If BB terminates with an empty \mathcal{R} , then stop (the point v^ν is an optimal solution of P).

When BB2 is used in the algorithm GOA, it can be infinite. For the case that Procedure BB2 is infinite, we introduce the following concept of approximate optimal solutions of Problem P .

DEFINITION 5.1 *Given a real number $\gamma > 0$, a flow x is called a γ -optimal solution of Problem P if it satisfies the following conditions:*

- (i) *There exists $\lambda \in \Lambda$ such that $\lambda x - \phi(\lambda) \geq -\gamma$, and*
- (ii) *dx is a lower bound of the optimal value of Problem P .*

Using this concept, we modify Procedure BB2 slightly to obtain the finiteness of the global optimization algorithm. Recall that $\lambda_k^1, \dots, \lambda_k^n$ are the vertices of S_k and (x_k^1, \dots, x_k^n) is an optimal solution of Problem $\bar{P}(S_k)$. The modification consists of the following additional stopping criterion between (a) and (b) at step $\langle k1 \rangle$:

$$\text{if } \sum_{\ell=1}^n \lambda_k^\ell x_k^\ell - \Phi(S_k) \geq -\delta, \quad \text{then stop.} \quad (5.1)$$

We will show in Theorem 5.2 below that $x_k = \sum_{\ell=1}^n x_k^\ell$ is an approximate optimal solution of Problem P in the sense of Definition 5.1.

THEOREM 5.2 *Assume that within Procedure BB2 the simplicial division is exhaustive, and the additional stopping criterion (5.1) is used at Step $\langle k1 \rangle$. Then*

- (i) *The global optimization algorithm always terminates after finitely many iterations; and*
- (ii) *If Procedure BB2 terminates by the additional stopping criterion (5.1), then $x_k = \sum_{\ell=1}^n x_k^\ell$ is a $(\delta + n\varepsilon)$ -optimal solution of Problem P , where*

$$\varepsilon = \max\{(\lambda_k^\ell - \lambda_k^{\ell'})x_k^\ell \mid \ell, \ell' = 1, \dots, n\}. \quad (5.2)$$

Proof We show that Procedure BB2 is finite. Then the finiteness of GOA follows immediately.

Suppose Procedure BB2 could be infinite. Then from Theorem 4.2, it would generate an infinite sequence $\{x_k\}$ converging to an optimal solution of Problem P . This implies that there exists an index \bar{k} such that

$$\sum_{\ell=1}^n \lambda_{\bar{k}}^{\ell} x_{\bar{k}}^{\ell} - \Phi(S_{\bar{k}}) \geq -\delta,$$

Thus, Procedure BB2 must stop at iteration \bar{k} .

Assume now Procedure BB2 terminates at step $\langle k1 \rangle$ of Iteration k by stopping criterion (5.1). From (5.2) it follows that for each $\ell = 1, \dots, n$ and for any $\lambda \in S_k$ we have

$$\lambda_k^{\ell} x_k^{\ell} - \lambda x_k^{\ell} \leq \varepsilon,$$

which implies

$$\lambda \sum_{\ell=1}^n x_k^{\ell} \geq \sum_{\ell=1}^n \lambda_k^{\ell} x_k^{\ell} - n\varepsilon.$$

Let now $\lambda' \in S_k$ such that $\phi(\lambda') = \Phi(S_k)$. Then

$$\lambda' x_k - \phi(\lambda') = \lambda' \sum_{\ell=1}^n x_k^{\ell} - \phi(\lambda') \geq \sum_{\ell=1}^n \lambda_k^{\ell} x_k^{\ell} - \Phi(S_k) - n\varepsilon,$$

which implies by (5.1) that

$$\lambda' x_k - \phi(\lambda') \geq -(\delta + n\varepsilon).$$

Note that $dx_k = \bar{\mu}(S_k)$ is a lower bound of the optimal value of P by the choice of S_k . Then by Definition 5.1 x_k is a $(\delta + n\varepsilon)$ -optimal solution of Problem P . ■

6 AN ILLUSTRATIVE EXAMPLE AND CONCLUSIONS

To illustrate the global optimization algorithm GOA we consider the network with 4 + 2 nodes and 10 arcs shown in Fig. 1, where the number attached to each arc is the arc capacity c_1, \dots, c_{10} .




FIGURE 1 Network with 6 nodes and 10 arcs.

For this example, Procedure BB1 is chosen for checking up global optimality. Within BB1 a lower bound for each rectangle S is computed by solving Problem $\bar{P}''(S, V)$ in Lemma 3.9, and the integral rectangular bisection is performed by the following rule.

Let $(x(S), \lambda(S), t(S))$ be an optimal solution of Problem $\bar{P}''(S, V)$. The rectangle S is then divided into $\{S_1, S_2\}$ by the integral rectangular bisection using $\lambda(S)$ and index ℓ chosen by $\lambda_\ell(S) = \max\{\lambda_i(S) \mid \lambda_i(S) < \bar{\lambda}_i, i = 1, \dots, n\}$.

$\langle\langle$ Initialization $\rangle\rangle$

An extreme maximal flow is computed: $w^0 = (7, 3, 1, 4, 2, 1, 6, 1, 2, 8)$.

$\langle\langle$ Iteration $\nu = 1$ $\rangle\rangle$

$\langle\nu 1\rangle$ Applying AVS we obtain a local optimal solution $v^1 = (7, 3, 0, 4, 2, 1, 7, 0, 2, 8)$. Set $\alpha_1 = dv^1 - 1 = 9$.

$\langle\nu 2\rangle$ BB1 starts with the rectangle $S_0 = \{\lambda \in \mathbb{R}_{10} \mid (1, \dots, 1) \leq \lambda \leq (10, \dots, 10)\}$ and the set $V = \{w^0, v^1\}$. Solving $\bar{P}''(S_0, V)$ we obtain

$$\begin{aligned} \mu(S_0) &= 7.2706, \\ x(S_0) &= (4.2706, 3.0, 0.5591, 4.0, 0.2706, 0.0, 6.7115, 0.2885, 0.0, 7.2706), \\ \lambda(S_0) &= (6.4908, 10.0, 6.0322, 10.0, 2.2177, 1.0, 9.6290, 3.5968, 1.0, 9.1794). \end{aligned}$$

The solution $x(S_0)$ is not a maximal flow. We then divide S_0 into two subrectangles using $\lambda(S_0)$ and the index $\ell = 7$ chosen by the above rule.

At iteration 21 while handling the rectangle $S = \{\lambda \in \mathbb{R}_{10} \mid (6, 1, 8, 1, 1, 1, 10, 1, 1, 1) \leq \lambda \leq (7, 10, 8, 10, 10, 10, 10, 10, 5)\}$, we obtain by solving Problem $\bar{P}''(S, V)$

$$\begin{aligned} \bar{\mu}(S) &= 9.0, \\ x(S) &= (6.0, 3.0, 1.0, 4.0, 2.0, 0.0, 7.0, 0.0, 1.0, 8.0), \\ \lambda(S) &= (6.8571, 10.0, 8.0, 10.0, 10.0, 1.0, 10.0, 2.0, 5.5, 5.0). \end{aligned}$$

The point $x(S)$ is a maximal flow with $dx(S) = 9 = \alpha_1$. So, we set $w_2 = x(S)$ and go to iteration $\nu = 2$. Procedure BB1 stops here with the set \mathcal{R}_{21} consisting of 21 rectangles.




FIGURE 2 A minimum maximal flow (x_i^*, c_A) .

$\langle\langle$ Iteration $v = 2$ $\rangle\rangle$

$\langle v1$ Starting at w^1 apply AVS to obtain a local optimal solution $v^2 = w^1$. Set $\alpha_2 = dv^2 = 9 - 1 = 8$.

$\langle v2$ Procedure BB1 continues with \mathcal{R}_{21} and $\alpha_2 = 8$. BB1 terminates at iteration 893 with $\mathcal{R}_{893} = \emptyset$.

A global optimal solution $x^* = (6, 3, 1, 4, 2, 0, 7, 0, 1, 8)$ thus obtained with optimal value 9 is shown in Fig. 2, where (x_i^*, c_i) is given on each arc.

In this article we have combined different techniques in local search and global optimization to propose the algorithm for solving the minimum maximal flow problem. The characteristic property of this algorithm is that the advantageous network flow structure is fully exploited. A detailed implementation and comparison of different procedures are left for future research.

Acknowledgments

The authors would like to thank Dolf Talman, Tilburg University and Maiko Shigeno, University of Tsukuba for stimulating discussion. The first author is supported by the Grant-in-Aid for Young Scientists (B) 14780343, and the second and third authors are supported by the Grant-in-Aid for Scientific Research (B)(2) 14380188 of the Ministry of Education, Culture, Sports, Science and Technology of Japan.

References

- [1] P.K. Ahuja, T.L. Magnanti and J.B. Orlin (1993). *Network Flows, Theory, Algorithms, and Applications*. Prentice Hall, Englewood Cliffs.
- [2] H.A. Lethi, D.T. Pham and L.D. Muu (1996). Numerical solution for optimization over the efficient set by d.c. optimization algorithms. *Operations Research Letters*, **19**, 117–128.
- [3] K. Belling-Seib, P. Mevet and C. Muller (1988). Network flow problems with one side constraint: a comparison of three solution methods. *Computers and Operations Research*, **15**, 381–394.
- [4] H.P. Benson (1991). An all-linear programming relaxation algorithm for optimizing over the efficient set. *Journal of Global Optimization*, **1**, 83–104.
- [5] H.P. Benson (1992). A finite nonadjacent extreme-point search algorithm for optimization over the efficient set. *Journal of Optimization Theory and Applications*, **73**, 47–64.
- [6] H.P. Benson (1995). A geometric analysis of the efficient outcome set in multiple objective convex program with linear criteria functions. *Journal of Global Optimization*, **6**, 213–251.
- [7] H.P. Benson and D. Lee (1996). Outcome-based algorithm for optimizing over the efficient set of a bicriteria linear programming. *Journal of Optimization Theory and Applications*, **88**, 77–105.
- [8] H.P. Benson and S. Sayin (1994). Optimization over the efficient set: four special case. *Journal of Optimization Theory and Applications*, **80**, 3–18.
- [9] S. Bolintineanu (1993). Minimization of a quasi-concave function over an efficient set. *Mathematical Programming*, **61**, 89–110.
- [10] N. Bryson (1991). Applications of the parametric programming procedure. *European Journal of Operational Research*, **54**, 66–73.
- [11] J.P. Dauer and T.A. Fosnaugh (1995). Optimization over the efficient set. *Journal of Global Optimization*, **7**, 261–277.
- [12] J.G. Ecker and J.H. Song (1994). Optimizing a linear function over an efficient set. *Journal of Optimization Theory and Applications*, **83**, 541–563.
- [13] J. Fülöp, (1994). A cutting plane algorithm for linear optimization over the efficient set. In: S. Komlősi, T. Rapsák and S. Shaible (Eds.), *Generalized Convexity*, Lecture Notes in Economics and Mathematical Systems 405, pp. 374–385, Springer-Verlag, Berlin.
- [14] M.R. Garay and D.S. Johnson (1979). *Computers and Intractability: A Guide to the Theory of NP-Completeness*. Freeman, San Francisco.
- [15] R. Horst, P. Pardalos and N.V. Thoai (2000). *Introduction to Global Optimization*, 2nd extended ed. Kluwer Academic Publishers, Dordrecht.

- [16] R. Horst and N.V. Thoai (1999). Maximizing a concave function over the efficient or weakly-efficient set. *European Journal of Operational Research*, **117**, 239–252.
- [17] R. Horst and H. Tuy (1996). *Global Optimization: Deterministic Approach*, 3rd, revised and enlarged ed. Springer, Berlin.
- [18] M. Iri (1994). An essay in the theory of uncontrollable flows and congestion. Technical Report, Department of Information and System Engineering, Faculty of Science and Engineering, Chuo University, TRISE 94–03.
- [19] M. Iri (1996). Network flow-theory and applications with practical impact. J. Dolezal and J. Fidler (Eds.), *System Modelling and Optimization*, pp. 24–36. Chapman & Hall, London.
- [20] L.D. Muu. A convex-concave programming method for optimizing over the efficient set. *Acta Mathematica Vietnamica*, **25**(1), 67–85.
- [21] P.H. Naccache (1978). Connectedness of the set of nondominated outcomes in multicriteria optimization. *Journal of Optimization Theory and Applications*, **25**, 459–467.
- [22] J. Philip (1972). Algorithms for the vector maximization problem. *Mathematical Programming*, **2**, 207–229.
- [23] T.Q. Phong and J.Q. Tuyen (2000). Bisection search algorithm for optimizing over the efficient set. *Vietnam Journal of Mathematics*, **28**, 217–226.
- [24] R.T. Rockafellar and R.J-B. Wets (1998). *Variational Analysis*. Springer-Verlag, Berlin.
- [25] S. Sayin (2000). Optimizing over the efficient set using a top-down search of faces. *Operations Research*, **48**, 65–72.
- [26] Y. Sawaragi, H. Nakayama and T. Tanino (1985). *Theory of Multiobjective Optimization*. Academic Press, Orland.
- [27] J.M. Shi and Y. Yamamoto (1997). A global optimization method for minimum maximal flow problem. *Acta Mathematica Vietnamica*, **22**, 271–287.
- [28] M. Shigeno, I. Takahashi and Y. Yamamoto (2003). Minimum maximal flow problem – An optimization over the efficient set. *Journal of Global Optimization*, **25**, 425–443.
- [29] R.E. Steuer (1985). *Multiple Criteria Optimization: Theory, Computation and Application*. Wiley, New York.
- [30] P.T. Thach, H. Konno and D. Yokota (1996). Dual approach to minimization on the set of Pareto-optimal solutions. *Journal of Optimization Theory and Applications*, **88**, 689–707.
- [31] N.V. Thoai and H. Tuy (1980). Convergent algorithms for minimizing a concave function. *Mathematics of Operations Research*, **5**, 556–566.
- [32] N.V. Thoai (2000). A class of optimization problems over the efficient set of a multiple criteria nonlinear programming problem. *European Journal of Operational Research*, **122**, 58–68.
- [33] N.V. Thoai (2000). Conical algorithm in global optimization for optimizing over efficient sets. *Journal of Global Optimization*, **18**, 321–336.
- [34] H. Tuy (1998). *Convex Analysis and Global Optimization*. Kluwer, Dordrecht.
- [35] D.J. White (1982). *Optimality and Efficiency*. John Wiley & Sons, Chichester.
- [36] D.J. White (1996). The maximization of a function over the efficient set via a penalty function approach. *European Journal of Operational Research*, **94**, 143–153.
- [37] S. Yamada, T. Tanino and M. Inuiguchi (2000). An inner approximation method for optimization over the weakly efficient set. *Journal of Global Optimization*, **16**, 197–217.
- [38] Y. Yamamoto (2002). Optimization over the efficient set: overview. *Journal of Global Optimization*, **22**, 285–317.